

1. From class we know that in 2D potential flow the virtual or apparent mass matrix for a body is given by

$$M_{ij} \equiv 2\pi\rho m_{ij} - \rho A_b \delta_{ij}$$

where A_b is the area of the body and the behavior at infinity of the potential is

$$\phi = -m_{ij}U_j x_i / |\mathbf{x}|^2 + o(|\mathbf{x}|^{-1}).$$

From potential flow past a flat plate calculate the apparent mass of a flat plate of width $4a$ moving normal to its surface. (Hint: Consider the steady flow past the plate, then obtain the disturbance potential from that.)

2. Consider a Navier-Stokes fluid of constant ρ, μ , no body forces. Consider a motion in a fixed bounded domain V with no-slip condition on its rigid boundary. Show that

$$dE/dt = -\Phi, E = \int_V \rho |\mathbf{u}|^2 / 2 dV, \Phi = \mu \int_V (\nabla \times \mathbf{u})^2 dV.$$


This shows that for such a fluid kinetic energy is converted into heat at a rate $\Phi(t)$. This last function of time gives the net *viscous dissipation* for the fluid contained in V . (Hint: $\nabla \times (\nabla \times \mathbf{u}) = \nabla(\nabla \cdot \mathbf{u}) - \nabla^2 \mathbf{u}$. Also $\nabla \cdot (\mathbf{A} \times \mathbf{B}) = \nabla \times \mathbf{A} \cdot \mathbf{B} - \nabla \times \mathbf{B} \cdot \mathbf{A}$.)

3. In two dimensions, with streamfunction ψ , where $(u, v) = (\psi_y, -\psi_x)$, show that the incompressible Navier-Stokes equations without body forces for a fluid of constant ρ, μ reduce to

$$\frac{\partial}{\partial t} \nabla^2 \psi - \frac{\partial(\psi, \nabla^2 \psi)}{\partial(x, y)} - \nu \nabla^4 \psi = 0.$$

In terms of ψ , what are the boundary conditions on a rigid boundary if the no-slip condition is satisfied there?

4. Consider the steady 2D flow of a layer of viscous incompressible fluid under gravity down an inclined plane. You may assume the streamlines are parallel to the plane, and that $(u, v) = (u(y), 0)$, where the x -axis is parallel to the plane (see the figure). Write down the equations for the flow, assuming constant ρ, μ . Solve for the pressure and for u , requiring that $p = p_0 = \text{constant}$ and $\mu du/dy = 0$ at the free surface adjacent to the air. (The latter condition imposes zero stress at the free surface). Compute the volume flux of fluid down the plane as a function of $\nu = \mu/\rho$, gravity g , and the layer thickness H .



5. Find the time-periodic 2D flow in a channel $-H < y < H$, filled with viscous incompressible fluid, given that the pressure gradient is $dp/dx = A + B \cos(\omega t)$, where A, B, ω are constants. This is an oscillating 2D Poiseuille flow. You may assume that $u(y, t)$ is even in y and periodic in t with period $2\pi/\omega$.