

# Calculus of Variations, Lecture 4, 2/19/2013

New topic: 1D problems

$$\min \int_a^b F(t, u(t), \dot{u}(t)) dt \quad u: [a, b] \rightarrow \mathbb{R}^n$$

emphasizing

- a) geodesics, as a key example
- b) importance of  $F$  being convex wrt  $\dot{u}$
- c) role of 2nd variation; conjugate pts

[Students taking Mechanics will see additional examples there, associated with solving eqns of Hamiltonian mechanics by "action minimization".]

Reasonable source for most of this material:  
Jost + Li-Jost, Sections 1.1-1.3 and 2.1.

Key example: geodesics. By defn: a geodesic is a curve that (locally) minimizes arc length.  
In local coordinates, if the curve is  $\bar{x}(t)$ ,

$$|\bar{x}| = |\dot{\bar{x}}(t)| dt = \left( \sum g_{ij}(\bar{x}(t)) \dot{x}_i \dot{x}_j \right)^{1/2}$$

where  $g_{ij}$  is the Riemannian metric; assoc var'l problem is

$$L = \int_a^b |\dot{x}(t)| dt$$

(note: we're interested in critical pts, not just minima).

Two issues:

(1) this has the form  $\int F(x(t), \dot{x}(t)) dt$  but  $F$  is not smooth in  $\dot{x}$  near  $\dot{x} = 0$

(2) arc length is indep of parametrization, so var'l pbm chooses a "curve" but not any particular parametrization (thus: a dramatic but geometrically-natural failure of uniqueness)

Both issues can be fixed by considering instead the different functional

$$E = \frac{1}{2} \int_a^b |\dot{x}|^2 dt$$

where  $|\dot{x}|^2 = \sum g_{ij}(x(t)) \dot{x}_i(t) \dot{x}_j(t)$ . To see why, observe that for any parametrized curve  $\tilde{x}(t)$ ,

$$L[x] \leq \sqrt{2(b-a)} \sqrt{E[x]}$$

(with strict inequality unless  $|\dot{x}|$  is constant), as a consequence of

$$\int_a^b |\dot{x}| dt \leq \left( \int_a^b |\dot{x}|^2 dt \right)^{1/2} \left( \int_a^b 1 dt \right)^{1/2}$$

Thus

$$\text{min value of } L \leq \sqrt{2(b-a)} \cdot (\text{min value of } E)^{\frac{1}{2}}.$$

But opposite  $\neq$  is easy: given any curve with length  $\ell$ , its constant-speed parametrization has  $|\dot{x}| = \ell/(b-a)$ , so

$$\frac{1}{2} \int_a^b |\dot{x}|^2 dt = \frac{1}{2}(b-a) \frac{\ell^2}{(b-a)^2} = \frac{1}{2(b-a)} \ell^2$$

Thus

$$(\text{min value of } E)^{\frac{1}{2}} \leq \frac{1}{\sqrt{2(b-a)}} (\text{min value of } L)$$

Conclusion: minimizers of  $E$  has min length and constant speed.

(Exercise: use the EL eqn for  $E$  to give a different proof that extremals - even critical pts! - of  $E$  have constant speed, by showing that  $\frac{d}{dt} |\dot{x}|^2 = 0$  if  $x(t)$  solves the EL eqn.).

Key properties of geodesics:

- a) They're smooth
- b) They're locally paths of shortest length
- c) globally, they may not be paths of shortest length (eg on a sphere the geodesics are arcs of great circles)

Rule: discuss above assumed we had a single "coordinate chart" valid along entire curve.  
 Locally true, but not necessarily globally so.  
 In general must use different coord charts on different parts of curve (see 3.2.1 of Post/Li-Post for detail on what this means).

Properties (a) - (c) are not special to geodesics; so it's natural to discuss them more generally, for problems of form

$$\int_a^b F(t, u(t), \dot{u}(t)) dt$$

where  $u: [a, b] \rightarrow \mathbb{R}^n$ . Note that EL eqn in this setting is

$$\frac{\partial F}{\partial u_j} - \frac{d}{dt} \frac{\partial F}{\partial \dot{u}_j} = 0 \quad 1 \leq j \leq n.$$

Discussion of (a) = smoothness of solns: we clearly need some condition on  $F$ , since for  $u: [-1, 1] \rightarrow \mathbb{R}$ ,

$$\min_{\substack{u(-1)=0 \\ u(1)=0}} \int_{-1}^{+1} (u_t^2 - 1)^2 dt \text{ is solved by } \begin{cases} u(t) = 0 & t \in [-1, 1] \\ u(-1) = 0 \\ u(1) = 0 \end{cases}$$



$$\min_{\substack{u(-1)=0 \\ u(1)=1}} \int_{-1}^{+1} (u_t^2 - 1)^2 dt \text{ is solved by } \begin{cases} u(t) = 0 & t \in [-1, 0] \\ u(t) = 2t & t \in [0, 1] \\ u(-1) = 0 \\ u(1) = 1 \end{cases}$$



Convenient hypothesis is that  $F(t, u, p)$  is smooth enough (I won't try to give minimal conditions - see fast + Li-fast for such things) and strictly convex in  $p$ . The point: the EL eqn can be written as

$$\frac{\partial F}{\partial \dot{u}_j} - \frac{\partial^2 F}{\partial u_j \partial t} - \sum_k \frac{\partial^2 F}{\partial \dot{u}_j \partial u_k} \ddot{u}_k = \sum_k \frac{\partial^2 F}{\partial \dot{u}_j \partial \dot{u}_k} \ddot{u}_k,$$

which we can solve (inverting the strictly pos. def. matrix  $\frac{\partial^2 F}{\partial \dot{u}_j \partial \dot{u}_k}$ ) to see that  $\ddot{u}$  is bounded if  $\dot{u}$  is bounded. Higher orders can be handled similarly (differentiate eqn in  $t$ ).

Proceeding argt is a bit sloppy, since it assumes  $\ddot{u}$  exists. Let's explain why strict convexity  $\Rightarrow$  it must exist. Consider

$$\varphi_j(t, u, p, g) = \frac{\partial F}{\partial p_j} - g_j$$

and observe that  $\vec{p}$  solves  $\vec{\varphi}(t, u, p, g) = 0$  iff it achieves

$$\max_p \langle g, p \rangle - F(t, u, p).$$

(Here  $t, u, g$  enter only as parameters; maximizing  $p$  is unique if  $F$  is strictly convex; argt has an implicit hypothesis that  $F(t, u, p)$  grows faster than

linearly as  $|p_j| \rightarrow \infty$ , so optimal  $p$  in preceding formula exists [ $p \rightarrow \infty$  is not optimal].)

Implicit function + hypothesis that  $\frac{\partial^2 F}{\partial p_i \partial p_k}$  has full rank  $\Rightarrow$  we can (locally) solve eqn  $\vec{q} = 0$  for  $p$  as fn of other vars, say

$$\frac{\partial F}{\partial p_j} = g_j \quad \forall j \iff p_j = \Psi_j(t, u, \vec{g}).$$

Now, we know  $\vec{q} = 0$  when  $\vec{g} = \frac{\partial F}{\partial p}$ . Evaluating this at  $p = \bar{u}$  gives

$$\dot{u}(t) = \Psi_j(t, u(t), \frac{\partial F}{\partial u}(t, u, \bar{u}))$$

RHS is diff'ble (using EL eqn to know differentiability of  $\frac{\partial F}{\partial u}$ )  $\Rightarrow$  LHS is diff'ble (in  $t$ ).

---

Rest of these notes discusses pts (b) + (c)  
(local minimality, conjugate pts, etc).

Brief summary :

- 1) 2<sup>nd</sup> variation provides a convenient necessary condition for minimality
- 2) importance of convexity is visible here too: if  $F$  is not convex in  $p$ , then 2<sup>nd</sup> var test is sure to fail.

3) as we work on longer time intervals  
 (eg  $[a, b]$ , with a fixed  $a$  &  $b \uparrow$ ) failure  
 of local optimality can be detected by  
 2nd varn test

(Suffit condls for optimality are also interesting of  
 course, but they would lead us too far astray.)

Defn of 2nd varn: given a soln  $u(t)$  of the EL eqn,  
 it is natural to consider

$$\frac{d^2}{ds^2} \Big|_{s=0} \int_a^b F(t, u+s\dot{\gamma}, \dot{u}+s\dot{\gamma}) dt$$

where  $\bar{\gamma}(t)$  is arbitrary (except perhaps for restrictions  
 to bdry condls). This reduces to

$$Q[\gamma] = \int_a^b F_{uu} \gamma \otimes \gamma + 2F_{u\dot{\gamma}} \gamma \otimes \dot{\gamma} + F_{\dot{\gamma}\dot{\gamma}} \dot{\gamma} \otimes \dot{\gamma} dt$$

where, for example,

$$F_{uu} \gamma \otimes \gamma = \sum \frac{\partial^2 F}{\partial u_i \partial u_j}(t, u, \dot{u}) \gamma_i(t) \gamma_j(t).$$

Focusing on case when  $u(a) + u(b)$  are fixed  
 (so  $\gamma(a) = \gamma(b) = 0$ ) we see that

$$u \text{ is loc min} \Rightarrow Q[\gamma] \geq 0 \text{ for all } \gamma \text{ s.t. } \gamma(a) = \gamma(b) = 0.$$

Importance of convexity is 2-fold.

First: If  $F_{pp} \geq c_0 I$  with  $c_0 > 0$  (this is a little stronger than strict convexity) Then

$$F_{pp}\dot{\gamma} \otimes \dot{\gamma} \geq c_0 |\dot{\gamma}|^2$$

and it's easy to see that  $Q$  is strictly positive if  $|b-a|$  is small enough. (Hint:  $\int_a^b |\dot{\gamma}|^2 \leq C(b-a)^2 \int_a^b |\ddot{\gamma}|^2$  if  $\gamma(a) = \gamma(b) = 0$ , with  $C$  only of  $b-a$ .)

Second: If, for some  $t_0$ , the matrix

$$\frac{\partial^2 F}{\partial p_i \partial p_j}(t_0, u(t_0), \dot{u}(t_0))$$


is not  $\geq 0$  (roughly:  $F$  is not convex in  $p$  at some point along the curve) then  $\exists \dot{\gamma} \neq 0$  s.t.  $Q[\dot{\gamma}] < 0$ . In fact, it suffices to choose  $\xi \in \mathbb{R}^n$  s.t.

$$\sum \frac{\partial^2 F}{\partial p_i \partial p_j}(t_0, u(t_0), \dot{u}(t_0)) \xi_i \xi_j < 0$$

and then take  $\dot{\gamma}(t)$  supported in a (small) nbhd of  $t_0$  s.t.

$$\dot{\gamma}(t) \in \{0, \pm \xi\}$$

Picture:



$$\dot{\gamma} = \begin{cases} \xi & t_0 - \varepsilon < t < t_0 \\ -\xi & t_0 < t < t_0 + \varepsilon \end{cases}$$

If  $\epsilon$  is small enough then  $Q[\gamma]$  is strictly negative (since  $\int_{\mathbb{P}} \dot{\gamma} \otimes \dot{\gamma} dt$  scales like  $\epsilon$  and is negative, while the other terms in  $Q$  scale like  $\epsilon^2$ ).

On long "thin intervals" local optimality can be lost.

Recall example of geodesics on a sphere!

We can detect loss of optimality using the 2nd variational quadratic form.

Observe that it makes sense to ask: does EL

$$\min_{\begin{array}{l} \gamma \\ \gamma(a) = 0 \\ \gamma(b) = 0 \end{array}} \int_a^b F_{uu} \dot{\gamma} \otimes \dot{\gamma} + 2F_{up} \dot{\gamma} \otimes \dot{\gamma} + F_{pp} \dot{\gamma} \otimes \dot{\gamma} dt$$

have a non-zero soln  $\gamma(t)$ ? (Such  $\gamma(t)$  solves the "homogeneous" 2nd order ODE

$$F_{uu} \ddot{\gamma} + F_{up} \dot{\gamma} - \frac{d}{dt}(F_{up} \dot{\gamma}) - \frac{d}{dt}(F_{pp} \dot{\gamma}) = 0$$

and is called a "Jacobi field".) If such  $\gamma(t)$  exists, we say  $b$  is conjugate to  $a$ .

Thm: After 1st conjugate pt, an extremal (ie a soln of EL eqns) ceases to be a minimizer.

Pf: Let  $b_0$  be conjugate to  $a$ , and  $b > b_0$ .

Try

$$(*) \quad \gamma = \begin{cases} \text{nonzero Jacobi field on } (a, b_0) \\ 0 \quad \text{on } (b_0, b) \end{cases}$$

I claim that 2nd varn evaluated at this  $\gamma$  vanishes.

Accepting this for a moment, the rest is easy:  
 The  $\gamma$  just defined is not  $C^2$ , so it cannot minimize the 2nd varn func (note: we have assumed that  $F_{pp} > 0$  so the term  $F_{pp}\dot{\gamma}\otimes\dot{\gamma}$  in the 2nd varn quad form is strictly convex). Thus I some other  $\tilde{\gamma}(t)$  for which 2nd varn is negative.

To see why 2nd varn eval at (\*) vanishes, consider

$$\Psi(t, \gamma, \dot{\gamma}) = F_{uu}\gamma \otimes \gamma + 2F_{up}\gamma \otimes \dot{\gamma} + F_{pp}\dot{\gamma} \otimes \dot{\gamma}$$

and observe that

$$\Psi(t, \alpha\gamma, \alpha\dot{\gamma}) = \alpha^2 \Psi(t, \gamma, \dot{\gamma}).$$

So (by diffn wrt  $\alpha$  at  $\alpha=1$ )

$$\gamma \cdot \Psi_\gamma + \dot{\gamma} \cdot \Psi_{\dot{\gamma}} = 2\Psi$$

Integrating :

$$\begin{aligned}\int_a^b \varphi(t, \gamma, \dot{\gamma}) dt &= \int_a^{b_0} \varphi(t, \gamma, \dot{\gamma}) dt \\ &= \frac{1}{2} \int_a^{b_0} \gamma \varphi_{\gamma} + \dot{\gamma} \varphi_{\dot{\gamma}} dt \\ &= \frac{1}{2} \int_a^{b_0} \gamma \left( \varphi_{\gamma} - \frac{d}{dt} \varphi_{\dot{\gamma}} \right) dt\end{aligned}$$

using that  $\gamma(a) = \gamma(b_0) = 0$ . But restriction of  $\gamma$  to  $[a, b_0]$  minimizes the 2nd order quadratic form on  $[a, b_0]$  (giving min value 0'), so it solves the EL eqns

$$\varphi_{\dot{\gamma}} - \frac{d}{dt}(\varphi_{\dot{\gamma}}) = 0 \quad \text{on } [a, b_0].$$

Therefore

$$\int_a^b \varphi(t, \gamma, \dot{\gamma}) dt = 0$$

as asserted.

I only proved in these notes that 2nd varn  $> 0$  on short intervals. But it's also true that crit pts are minimizers on short intervals if  $F_{PP}$  is pos def.

For special case of geodesics on  $S^2 \subset \mathbb{R}^3$ ,

The antipodal pt is the conjugate pt. To see why, observe that

- a) for antipodal pts, there is a 1-par family of shortest paths (great semicircles)
- b) if there's a 1-par family of minimizers  $u^\theta(t)$  then  $\eta = \frac{d}{d\theta} u^\theta$  is a Jacobi field.

Point (a) is obvious. For (b): each  $u^\theta$  solves EL eqn

$$F_u(t, u^\theta, \dot{u}^\theta) = \frac{d}{dt} F_p(t, u^\theta, \dot{u}^\theta).$$

Diffr wrt to  $\theta \Rightarrow$

$$F_{u\eta}(t, \eta + F_{up}\dot{\eta}) = \frac{d}{dt} (F_{p\eta}\eta + F_{p\dot{\eta}}\dot{\eta})$$

with  $\eta = \frac{d}{d\theta} u^\theta$ . This is precisely the eqn characterizing a Jacobi field (note that  $\eta=0$  at endpoints, i.e. poles).

[More conceptual pt: value of  $\int_a^b F(t, u^\theta, \dot{u}^\theta) dt$  wrt  $\theta \Rightarrow$  both 1st and 2nd derivs vanish wrt  $\theta$ . So (assuming  $u^\theta$  is extremal)  $\eta = \frac{d}{d\theta} u^\theta$  achieves value 0 in the 2nd variational. Since the min value is zero, this  $\eta$  must be a Jacobi field.]

(1) Show directly (using the EL eqns) that any extremal for  $\int_a^b |\dot{x}|^2 dt$  has constant speed (see pp 2-3).

(2) Show that if  $b$  is conjugate to  $a$  then

$$\min_{\substack{\gamma(a)=a \\ \gamma(b)=b}} \int_a^b F_{uu} \dot{u}^2 + 2F_{up} \dot{u} \dot{p} + F_{pp} \dot{p}^2 dt = 0$$

(3) When studying waves it is useful to consider paths that minimize "travel time," where the wave speed  $\gamma(x)$  is a known function of location  $x$ . Show that this amounts to considering geodesics in the metric

$$g_{ij} = \frac{1}{\gamma(x)} \delta_{ij}.$$

(4) In these notes we focused on Dir BC, i.e.  $\min \int_a^b F(t, u, \dot{u}) dt$  subject to  $u(a) = \alpha$  and  $u(b) = \beta$  being given. Suppose instead we impose  $u(a) = \alpha$  and  $u(b) \in M$  where  $M$  is a submanifold. What end condition does the EL get at  $t=b$ ? What is the proper notion of Jacobi field in this case?

(5) Show that the only critical pts of

$$\int_a^b u_x^2 + (u^2 - 1)^2$$

(no boundary condition!) with nonnegative 2nd variation are the "trivial ones," namely  $u \equiv -1$  and  $u \equiv +1$ . (Hint: let  $\varphi$  be a critical point. Show that  $\varphi = u_x$  achieves value 0 in the 2nd variation quadratic form. Then argue that this  $\varphi$  can't be a minimizer of that quadratic form.)