

APPROXIMATION AT PLACES OF BAD REDUCTION FOR RATIONALLY CONNECTED VARIETIES

BRENDAN HASSETT AND YURI TSCHINKEL

ABSTRACT. This paper addresses weak approximation for rationally connected varieties defined over the function field of a curve, especially at places of bad reduction. Our approach entails analyzing the rational connectivity of the smooth locus of singular reductions of the variety. As an application, we prove weak approximation for cubic surfaces and Fano hypersurfaces of dimension at least three, with square-free discriminant.

CONTENTS

1. Introduction	1
2. Notions of approximation	4
3. Curves, combs, and deformations	6
4. Strong rational connectivity	7
5. Cubic surfaces	14
6. Higher-dimensional Fano hypersurfaces	18
References	20

1. INTRODUCTION

In number theory, many results and techniques rely on approximating adelic points by rational points. In this paper, we study geometric versions of these notions for rationally connected varieties over the function field of a curve. In this context, rational points correspond to sections of rationally-connected fibrations over the curve. We are looking for sections with prescribed jet data in finitely many fibers.

Let k be an algebraically closed field of characteristic zero, B a smooth curve over k with function field $F = k(B)$. Let \overline{B} be the smooth projective model of F and put $S := \overline{B} \setminus B$.

Date: November 21, 2007.

Theorem 1. *Let X be a smooth proper rationally connected variety over F , and $\pi : \mathcal{X} \rightarrow B$ a proper model of X . (A model of X is an algebraic space flat over B with generic fiber X .) Let \mathcal{X}^{sm} be the locus where π is smooth and $\mathcal{X}^\bullet \subset \mathcal{X}^{sm}$ be an open subset such that*

- (1) *there exists a section $s : B \rightarrow \mathcal{X}^\bullet$;*
- (2) *for each $b \in B$ and $x \in \mathcal{X}_b^\bullet$, there exists a rational curve $f : \mathbb{P}^1 \rightarrow \mathcal{X}_b^\bullet$ containing x and the generic point of \mathcal{X}_b^\bullet .*

Then sections of $\mathcal{X}^\bullet \rightarrow B$ satisfy approximation away from S (see Section 2).

We shall actually prove a stronger result, Theorem 15, that is applicable in positive characteristic. Rationally-connected fibrations over curves have sections by [8]. The existence of a section through a finite set of prescribed points is addressed [16] 2.13 and [15] IV.6.10.1. Weak approximation is known in fibers of good reduction [10], so we take simultaneous resolutions of singular fibers of \mathcal{X} whenever possible. (For example, for rational double points on surfaces, simultaneous resolution is possible provided the local monodromy is trivial [3, 4].) Consequently, when $\mathcal{X} \rightarrow B$ admits a simultaneous resolution over some étale neighborhood of b , we replace \mathcal{X} by this resolution. However, the resolved family may be an algebraic space, rather than a scheme, over B . This is why Theorem 1 is stated in this generality.

There are very few instances where weak approximation over function fields is known at all places: stably rational varieties; connected linear algebraic groups and their homogeneous spaces; homogeneous space fibrations over varieties that satisfy weak approximation, including conic bundles over rational varieties; and Del Pezzo surfaces of degree at least four [5]. Recently, the case of smooth hypersurfaces of degree d in \mathbb{P}^n with $d^2 \leq n$ has been resolved [6], as an application of the notion of ‘rational simply connectedness’. Even the case of cubic surfaces remains open, in general. Madore established weak approximation for cubic surfaces at places of good reduction [18]. His proof uses the abundance of distinct unirational parametrizations, and builds on ideas of Swinnerton-Dyer [21].

When is Theorem 1 applicable? Let X be a smooth projective rationally connected variety over $F = k(B)$, with B projective. There exists a regular proper model $\pi : \mathcal{X} \rightarrow B$, and any section $s : B \rightarrow \mathcal{X}$ is

contained in \mathcal{X}^{sm} . For each singular fiber \mathcal{X}_b , fix an irreducible component $\mathcal{X}_b^\bullet \subset \mathcal{X}_b^{sm}$; these determine an open subset $\mathcal{X}^\bullet \subset \mathcal{X}^{sm}$. To prove weak approximation for X , it suffices to prove approximation for each \mathcal{X}^\bullet obtained in this way. We do not know how to verify (1) in general: Is there *any* section meeting a prescribed irreducible component of \mathcal{X}_b^{sm} ? Further, there is no general result giving a regular proper model $\mathcal{X} \rightarrow B$ such that each irreducible component of \mathcal{X}_b^{sm} has the property (2).

Section 5 is devoted to applications to cubic surfaces:

Theorem 2. *Let X be a smooth cubic surface over F and $\pi : \mathcal{X} \rightarrow B$ a proper model whose singular fibers are cubic surfaces with rational double points. Suppose there exists a section $s : B \rightarrow \mathcal{X}^{sm}$. Then sections of $\mathcal{X}^{sm} \rightarrow B$ satisfy approximation away from S .*

When the model is regular all sections are contained in the smooth locus, so we conclude:

Corollary 3. *Let X be a smooth cubic surface over F . Suppose X admits a regular proper model $\pi : \mathcal{X} \rightarrow B$ whose singular fibers are cubic surfaces with rational double points. Then weak approximation holds for X away from $S = \overline{B} \setminus B$.*

There exist cubic surfaces which do not admit models with at most rational double points in a given fiber, e.g., the isotrivial family

$$x^3 + y^3 + z^3 = tw^3$$

over the t -line. Nonetheless, Corollary 3 proves weak approximation for ‘generic’ cubic surfaces.

Corollary 4. *Let $\mathcal{Hilb} = \mathbb{P}(\Gamma(\mathcal{O}_{\mathbb{P}^3}(3))) \simeq \mathbb{P}^{19}$ denote the Hilbert scheme of cubic surfaces, $\mathcal{U} \rightarrow \mathcal{Hilb}$ the universal family, and $D \subset \mathcal{Hilb}$ the discriminant divisor. Let $B \subset \mathcal{Hilb}$ be a smooth curve transverse to each branch of D and*

$$\mathcal{X} := \mathcal{U} \times_{\mathcal{Hilb}} B \rightarrow B$$

the corresponding family. Then $X = \mathcal{U} \times_{\mathcal{Hilb}} \text{Spec}(F)$ satisfies weak approximation away from $S = \overline{B} \setminus B$.

Note that meeting the discriminant transversally is an open condition on the classifying map to the Hilbert scheme. This can be expressed in number-theoretic terms: The discriminant of $\mathcal{X} \rightarrow B$ is square-free. Generalizations to degree-two del Pezzo surfaces have been proven by Knecht [12].

Finally, in Section 6 we offer extensions of these Corollaries. We generalize Corollary 3 to the case where the fibers have isolated complete-intersection terminal singularities. Corollary 4 extends to generic hypersurfaces $X \subset \mathbb{P}^n$ of degree $d \leq n$ provided $n \geq 4$.

In our approach to approximation, we require precise control over proper rational curves in the smooth locus of a singular variety. One focus of this paper is to extend standard results on smooth proper rationally connected varieties to the non-proper case (see Section 4). The application to cubic surfaces and higher-dimensional hypersurfaces involves refining rational connectivity results of [11] (see Sections 5 and 6).

Acknowledgments: We are grateful to J. L. Colliot-Thélène for numerous discussions about the problems considered here; the ideas here were developed during visits to Orsay by both authors. J. McKernan suggested the extensions to higher-dimensional hypersurfaces. We also benefitted from conversations with S. Keel, A. Knecht, and J. Kollar. The first author was partially supported by the Sloan Foundation and NSF Grants 0134259 and 0196187. The second author was partially supported by National Science Foundation Grants 0554280 and 0602333.

2. NOTIONS OF APPROXIMATION

Let F be a global field, i.e., a number field or the function field of a curve B defined over an algebraically closed field k . Let S a finite set of places of F containing the archimedean places, $\mathfrak{o}_{F,S}$ the corresponding ring of integers, and $\mathbb{A}_{F,S}$ the restricted direct product over all places outside S .

Let X be an algebraic variety over F , $X(F)$ the set of F -rational points and $X(\mathbb{A}_{F,S}) \subset \prod_{v \notin S} X(F_v)$ the set of $\mathbb{A}_{F,S}$ -points of X . The set $X(\mathbb{A}_{F,S})$ carries a natural direct product topology. One says that *weak approximation* holds for X away from S if $X(F)$ is dense in this topology.

The set $X(\mathbb{A}_{F,S})$ also carries a natural adelic topology: The basic open subsets are

$$\prod_{v \in S'} \mathfrak{u}_v \times \prod_{v \notin (S \cup S')} \mathcal{X}(\mathfrak{o}_v),$$

where S' is a finite set of nonarchimedean places disjoint from S , $\mathcal{X} \rightarrow \text{Spec}(\mathfrak{o}_{F,S})$ is a model of X (i.e., flat with generic fiber X), \mathfrak{o}_v is the completion of $\mathfrak{o}_{F,S}$ at v , and $\mathfrak{u}_v \subset X(F_v)$ an open subset in the v -adic

analytic topology on $X(F_v)$. This depends only on X not on the choice of model. *Strong approximation* holds for X away from S if $X(F)$ is dense in $X(\mathbb{A}_{F,S})$. Note that strong approximation implies weak approximation. Conversely, for $\mathcal{X} \rightarrow \text{Spec}(\mathfrak{o}_{F,S})$ flat and proper, weak approximation implies strong approximation, since $\mathcal{X}(\mathfrak{o}_v) = X(F_v)$; in these cases, we will use the term weak approximation for the sake of consistency.

Finally, there is a formulation which is sensitive to the choice of model. Consider the topology on $\prod_{v \notin S} \mathcal{X}(\mathfrak{o}_v)$ with basic open subsets

$$\prod_{v \in S'} \mathfrak{u}_v \times \prod_{v \notin (S \cup S')} \mathcal{X}(\mathfrak{o}_v),$$

with $\mathfrak{u}_v \subset \mathcal{X}(\mathfrak{o}_v)$ an open subset. We say that *approximation holds for S -integral points of \mathcal{X}* if $\mathcal{X}(\mathfrak{o}_{F,S})$ is dense in this product. This is a version of strong approximation *for integral points*.

We now focus on the function field case: Let \overline{B} be a smooth projective model of B with $S = \overline{B} \setminus B$; places v correspond to points $b \in \overline{B}$. Let X be a smooth variety proper over $F = k(B)$, $\pi : \mathcal{X} \rightarrow B$ a model proper and flat over B (which exists by [20]), and $\mathcal{X}^\bullet \subset \mathcal{X}^{sm}$ an open subset surjecting onto B . Since π is proper, F -rational points of X correspond to sections $s : B \rightarrow \mathcal{X}$. If \mathcal{X} is regular s factors through \mathcal{X}^{sm} .

Definition 5. An *admissible section* of $\pi : \mathcal{X} \rightarrow B$ is a section $s : B \rightarrow \mathcal{X}^{sm}$. An *admissible N -jet* of π at b is a section of

$$\mathcal{X}^{sm} \times_B \text{Spec}(\mathcal{O}_{B,b}/\mathfrak{m}_{B,b}^{N+1}) \rightarrow \text{Spec}(\mathcal{O}_{B,b}/\mathfrak{m}_{B,b}^{N+1}).$$

An *approximable N -jet* of π at b is a section of

$$\mathcal{X} \times_B \text{Spec}(\mathcal{O}_{B,b}/\mathfrak{m}_{B,b}^{N+1}) \rightarrow \text{Spec}(\mathcal{O}_{B,b}/\mathfrak{m}_{B,b}^{N+1})$$

that may be lifted to a section of $\widehat{\mathcal{X}}_b \rightarrow \widehat{B}_b$, with $\widehat{B}_b = \text{Spec}(\hat{\mathcal{O}}_{B,b})$ and $\widehat{\mathcal{X}}_b = X \times_B \widehat{B}_b$.

Hensel's lemma guarantees that every admissible N -jet is approximable. Let $\{b_i\}_{i \in I}$ be a finite set of points and j_i an admissible N -jet of π at b_i . We write $J = \{j_i\}_{i \in I}$ for the corresponding collection of admissible N -jets.

The notions of weak and strong approximation introduced above have geometric interpretations

- Weak and strong approximation hold for X away from S if any finite collection of approximable jets of π can be realized by a section $s : B \rightarrow \mathcal{X}$.
- This is equivalent to weak approximation holding for X^\bullet (the generic fiber of \mathcal{X}^\bullet) away from S : Every jet in \mathcal{X} at b can be realized by a section $\mathcal{X} \times_B \widehat{B}_b \rightarrow \widehat{B}_b$ meeting $\widehat{\mathcal{X}}_b^\bullet$.
- If \mathcal{X} is regular these are equivalent to the condition that any collection of admissible jets of π can be realized by a section $s : B \rightarrow \mathcal{X}^{sm}$.

There is an analogous formulation of approximation for integral points:

- Approximation holds for sections of $\mathcal{X}^\bullet \rightarrow B$ away from S if each collection of jet data in \mathcal{X}^\bullet can be realized by a section $s : B \rightarrow \mathcal{X}^\bullet$.
- If \mathcal{X} is regular and $\mathcal{X}^\bullet = \mathcal{X}^{sm}$ this is equivalent to weak approximation for X .

3. CURVES, COMBS, AND DEFORMATIONS

The *dual graph* associated with a nodal curve C has vertices indexed by the irreducible components of C and its edges indexed by the intersections of these components. A projective nodal curve C is *tree-like* if

- each irreducible component of C is smooth;
- the dual graph of C is a tree.

Definition 6. A *comb with m reducible teeth* is a projective nodal curve C with $m + 1$ subcurves D, T_1, \dots, T_m such that

- D is smooth and irreducible;
- $T_l \cap T_{l'} = \emptyset$, for all $l \neq l'$;
- each T_l meets D transversally in a single point; and
- each T_l is a chain of \mathbb{P}^1 's.

Here D is called the *handle* and the T_l the *reducible teeth*.

Let C be a nodal curve and $h : C \rightarrow W$ an immersion into a smooth algebraic space with nodal image. (In particular, h is an embedding at nodes of C .) Let \mathcal{N}_h denote the normal bundle (or sheaf), i.e., the dual to the kernel of the restriction $h^* \Omega_W^1 \rightarrow \Omega_C^1$.

We will use the following lemma, which has the same proof as Proposition 24 of [10]:

Lemma 7. *Let C be a tree-like curve, W a smooth algebraic space, $h : C \rightarrow W$ an immersion with nodal image. Suppose that for each irreducible component C_l of C , $H^1(C_l, \mathcal{N}_h \otimes \mathcal{O}_{C_l}) = 0$ and $\mathcal{N}_h \otimes \mathcal{O}_{C_l}$ is globally generated. Then h deforms to an immersion of a smooth curve into W .*

Suppose furthermore that $\mathfrak{w} = \{w_1, \dots, w_M\} \subset C$ is a collection of smooth points such that for each component C_l , $H^1(\mathcal{N}_h \otimes \mathcal{O}_{C_l}(-\mathfrak{w})) = 0$ and the sheaf $\mathcal{N}_h \otimes \mathcal{O}_{C_l}(-\mathfrak{w})$ admits a section nonvanishing at each point of the support of

$$(\mathcal{N}_h \otimes \mathcal{O}_{C_l})/\mathcal{N}_{h|C_l}.$$

Then $h : C \rightarrow W$ deforms to an immersion of a smooth curve into W containing $h(\mathfrak{w})$.

4. STRONG RATIONAL CONNECTIVITY

Definition 8. A variety X is *rationally connected* (resp. *separably rationally connected*) if there is a family of proper irreducible rational curves $g : U \rightarrow Z$ (resp. $\pi_2 : U = \mathbb{P}^1 \times Z \rightarrow Z$) and a cycle morphism $u : U \rightarrow X$ such that

$$u^2 : U \times_Z U \rightarrow X \times X$$

is dominant (resp. smooth over the generic point)).

Intuitively, two generic points of X can be joined by an irreducible projective rational curve. Over fields of characteristic zero, rational connected varieties are also separably rationally connected [15] IV.3.3.1.

The notion of rational connectedness is a bit subtle over countable fields [2]. For convenience, we work over an uncountable algebraically closed field. Over such a field, rational connectivity is equivalent to the condition that two very general points of X can be joined by such a rational curve.

Definition 9. Let X be a smooth algebraic space of dimension d and $f : \mathbb{P}^1 \rightarrow X$ a nonconstant morphism, so we have an isomorphism

$$f^* \mathcal{T}_X \simeq \mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \dots \oplus \mathcal{O}_{\mathbb{P}^1}(a_d)$$

for suitable integers a_1, \dots, a_d . Then f is *free* (resp. *very free*) if each $a_i \geq 0$ (resp. $a_i \geq 1$).

We refer the reader to [15] IV.3 for further facts about rationally connected varieties.

One technical result will play a prominent rôle in our analysis.

Proposition 10 ([15] IV.3.9.4). *Let V be a smooth separably rationally connected (not necessarily proper) variety. Then there exists a nonempty subset $V^0 \subset V$ characterized as the largest open subset such that if $v_1, \dots, v_m \in V^0$ are distinct closed points, then there is a very free curve in V^0 containing these as smooth points. Moreover, any rational curve $C \subset V$ that meets V^0 is contained in V^0 .*

No example where $V^0 \neq V$ is known.

Remark 11. Let V_2 be a smooth variety, $V_1 \subset V_2$ a rationally connected dense open subvariety, and $V_2^0 \subset V_2$ the largest open set satisfying the conditions of Proposition 10. Then $V_1^0 \subset V_2^0$. Thus a point $v \in V_2$ is in V_2^0 provided there is a rational curve $f : \mathbb{P}^1 \rightarrow V_2$ through v and meeting V_1^0 .

Proposition 12. *Let V be a smooth separably rationally connected variety, and $\beta : W \rightarrow V$ an iterated blow-up of V along smooth subvarieties. Then $\beta^{-1}(V^0) = W^0$.*

Proof. The inclusion $W^0 \subset \beta^{-1}(V^0)$ is straightforward: Given points $w_1, \dots, w_m \in W^0$, there is a very free curve $g : \mathbb{P}^1 \rightarrow W^0$ containing them; we may choose this to be transversal to the exceptional divisor of β . The inclusion of sheaves

$$\mathcal{T}_W \hookrightarrow \beta^* \mathcal{T}_V$$

remains an inclusion after pull-back via g , as the support of the cokernel does not contain $g(\mathbb{P}^1)$. The positivity of $g^* \mathcal{T}_W$ implies the positivity of $(\beta \circ g)^* \mathcal{T}_V$, which means that $\beta \circ g : \mathbb{P}^1 \rightarrow V$ is also very free.

For the reverse direction, we may restrict to the case where W is the blow-up of V along a smooth subvariety Z of codimension $r > 1$, with exceptional divisor E . It is clear that $\beta^{-1}(V^0 \setminus Z) \subset W_0$, so consider some $w \in \beta^{-1}(z)$ with $z \in Z \cap V^0$. It suffices to construct a rational curve containing w and the generic point of W .

There exists a very free curve $f' : \mathbb{P}^1 \rightarrow V^0$ with the following properties:

- (1) $f'(\mathbb{P}^1)$ meets Z only at z (we can always deform a very free curve to a curve passing through z and disjoint from a codimension ≥ 2 subset [15, II.3.7]);
- (2) $f'(\mathbb{P}^1)$ is smooth at z and transverse to Z .

Let $g' : \mathbb{P}^1 \rightarrow W$ denote the lift to W , which is free in W , and $w' = g'(0)$. If $w' = w$ then we are done. Otherwise, let $\ell \subset \beta^{-1}(z) \simeq \mathbb{P}^{r-1}$ denote the

line joining w and w' . Since g' is free, it admits a small deformation to a free curve $g'': \mathbb{P}^1 \rightarrow W$ with $w'':=g''(0) \in \ell, w'' \neq w'$. (See Figure 1.)




FIGURE 1. Constructing the comb

We construct a comb $h: C \rightarrow W$ with handle $\ell \subset \mathbb{P}^{r-1} \subset W$ and two teeth $g', g'': \mathbb{P}^1 \rightarrow W$. Since E is exceptional, we compute

$$\mathcal{N}_{\ell/E} \simeq \mathcal{O}_{\mathbb{P}^1}^{\dim(V)} \oplus \mathcal{O}_{\mathbb{P}^1}(1)^{r-2}$$

and

$$\mathcal{N}_{E/W} \otimes \mathcal{O}_\ell \simeq \mathcal{O}_{\mathbb{P}^1}(-1).$$

The exact sequence of normal bundles

$$0 \rightarrow \mathcal{N}_{\ell/E} \rightarrow \mathcal{N}_{\ell/W} \rightarrow \mathcal{N}_{E/W} \otimes \mathcal{O}_\ell \rightarrow 0$$

splits because

$$\text{Ext}^1(\mathcal{O}_{\mathbb{P}^1}(-1), \mathcal{O}_{\mathbb{P}^1}(n)) = 0, n \geq -2.$$

Thus we deduce

$$\mathcal{N}_{\ell/W} \simeq \mathcal{O}_{\mathbb{P}^1}^{\dim(V)-r} \oplus \mathcal{O}_{\mathbb{P}^1}(1)^{r-2} \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$$

where the negative summand is in the normal direction to E . Since $g'(\mathbb{P}^1)$ and $g''(\mathbb{P}^1)$ are transverse to E , we can apply Proposition 23 of [10]. The key point is that the only negativity in the normal bundle of ℓ is due to the negativity of the normal bundle of $E \subset W$; however, the components $g'(\mathbb{P}^1)$ and $g''(\mathbb{P}^1)$ overcome this. Precisely, we have

$$\mathcal{N}_h \otimes \mathcal{O}_\ell \simeq \mathcal{O}_{\mathbb{P}^1}^{\dim(V)-r} \oplus \mathcal{O}_{\mathbb{P}^1}(1)^{r-2} \oplus \mathcal{O}_{\mathbb{P}^1}(1);$$

the quotient $(\mathcal{N}_h \otimes \mathcal{O}_\ell)/\mathcal{N}_{\ell/W}$ lies in the image of the last summand.

Lemma 7 implies that $h: C \rightarrow W$ admits a deformation to a rational curve containing w . \square

A similar argument gives the following strengthening of Proposition 10 (cf. Theorem 2.2 of [7])

Proposition 13. *Let V be a smooth separably rationally connected variety and $V^0 \subset V$ be the distinguished open subset characterized in Proposition 10. Then for any finite collection of jets*

$$j_i : \text{Spec}(k[\epsilon]/\langle \epsilon^{N+1} \rangle) \hookrightarrow V^0, \quad i = 1, \dots, m$$

supported at distinct points v_1, \dots, v_m , there exists a very free rational curve smooth at v_1, \dots, v_m with the prescribed jets.

Proof. There is an iterated blow-up

$$\beta : W = W_N \rightarrow \dots \rightarrow W_j \rightarrow \dots \rightarrow W_1 \rightarrow V$$

and points $w_1, \dots, w_m \in W$ so that if $g : C \rightarrow W$ is a morphism whose image contains w_1, \dots, w_m then the image of $f := \beta \circ g : C \rightarrow V$ contains the given collection of jets. Here is the description: Over each point v_i , we blow up V successively at N points. Given any smooth curve germ C with the prescribed N -jet at v_i , W_j is the blowup of W_{j-1} at the points of the proper transform of C lying over the v_i . Proposition 12 then implies there exists a very free curve $g : \mathbb{P}^1 \rightarrow W$ through w_1, \dots, w_m . However, the image of this curve in V will be singular at v_i if $g(\mathbb{P}^1)$ meets $\beta^{-1}(v_i)$ in more than one point.

We claim there exists a very free curve $g_i : \mathbb{P}^1 \rightarrow W$ meeting $\beta^{-1}(v_i)$ only at w_i , transversally. We choose this curve so that it is disjoint from $\beta^{-1}(v_j)$ when $j \neq i$. (Again, we are using the fact that a very free curve can be deformed away from any codimension ≥ 2 subvariety while passing through a prescribed point in the complement.) Fix generic points $x_i \in g_i(\mathbb{P}^1)$ and let $g_0 : \mathbb{P}^1 \rightarrow W$ be a very free curve intersecting $g_i(\mathbb{P}^1)$ transversely at x_i but not meeting any $\beta^{-1}(v_i)$. (For example, take $g_0 = (\beta^{-1} \circ f_0)$, where $f_0 : \mathbb{P}^1 \rightarrow V$ is a very free curve through $\beta(x_1), \dots, \beta(x_m)$.) Consider the comb $h : C \rightarrow W$ with handle $g_0(\mathbb{P}^1)$ and m -teeth $g_i(\mathbb{P}^1)$. This deforms to a very free curve $h' : \mathbb{P}^1 \rightarrow W$ meeting each $\beta^{-1}(v_i)$ only at w_i , transversally.

The proof of the claim is a refinement of the argument for Proposition 12. We proceed by induction on N . The base case $N = 1$ is contained in the proof of Proposition 12, which gives a very free curve smooth at v_i with prescribed tangency. Let $E_{i,N} \simeq \mathbb{P}^{\dim(V)-1}$ be the last exceptional divisor of $\beta : W \rightarrow V$ over v_i , i.e., the exceptional divisor of the N -th blow-up. For $1 \leq j < N$, let $E_{i,j} \subset W_N$ denote the proper transform of the exceptional divisor of $W_j \rightarrow W_{j-1}$ over v_i ; we

have $E_{i,j} \simeq \text{Bl}_{w_{i,j}} \mathbb{P}^{\dim(V)-1}$, where $w_{i,j}$ is the intersection of the proper transform of C with the exceptional divisor of $W_j \rightarrow W_{j-1}$.

Suppose that $g'_i : \mathbb{P}^1 \rightarrow W$ is a very free curve such that $\beta \circ g'_i$ is smooth with the desired $(N-1)$ -jet at v_i . Let $w'_i = g'_i(\mathbb{P}^1) \cap \beta^{-1}(v_i)$ denote the unique point of intersection, which we assume is distinct from w_i . Let ℓ_N denote the line in $E_{i,N} \simeq \mathbb{P}^{\dim(V)-1}$ joining w_i and w'_i , and z_{N-1} its point of intersection with $E_{i,N-1}$. Let $\ell_{N-1} \subset E_{i,N-1} \simeq \text{Bl}_{w_{i,N-1}} \mathbb{P}^{\dim(V)-1}$ denote the proper transform of a line containing z_{N-1} , and z_{N-2} its point of intersection with $E_{i,N-2}$. Continue in this way, until we obtain $\ell_1 \subset E_{i,1}$, the proper transform of a line containing z_1 . Finally, let $g''_i : \mathbb{P}^1 \rightarrow W$ be a very free curve meeting the exceptional locus transversally at a generic point of ℓ_1 . (See Figure 2.)




FIGURE 2. Constructing the comb with reducible teeth

Let $h : C \rightarrow W$ be the comb with handle ℓ_N and two reducible teeth:

- (1) $g'_i : \mathbb{P}^1 \rightarrow W$;
- (2) the union of the lines $\ell_{N-1}, \dots, \ell_1$ and the curve $g''_i : \mathbb{P}^1 \rightarrow W$;

By a normal bundle computation similar to that of Proposition 12, we find that $\mathcal{N}_h|_{\ell_N}$ is ample and \mathcal{N}_h is nonnegative on each of the remaining components: Again, Lemma 7 (or Proposition 24 of [10]) implies that h admits a deformation to an immersed rational curve containing w_i .

Here are the details of the computations (cf. [10] Section 5): The normal bundle of a line in projective space is

$$\mathcal{N}_{\ell_N/E_{i,N}} = \mathcal{N}_{\ell_N/\mathbb{P}^{\dim(V)-1}} \simeq \mathcal{O}_{\mathbb{P}^1}(1)^{\dim(V)-2}$$

and the normal bundle for an exceptional divisor is

$$\mathcal{N}_{E_{i,N}/W} \simeq \mathcal{O}_{\mathbb{P}^{\dim(V)-1}}(-1).$$

For each j we have

$$(4.1) \quad 0 \rightarrow \mathcal{N}_{\ell_j/E_{i,j}} \rightarrow \mathcal{N}_{\ell_j/W} \rightarrow \mathcal{N}_{E_{i,j}/W}|_{\ell_j} \rightarrow 0$$

which for $j = N$ yields

$$\mathcal{N}_{\ell_N/W} \simeq \mathcal{O}_{\mathbb{P}^1}(1)^{\dim(V)-2} \oplus \mathcal{O}_{\mathbb{P}^1}(-1),$$

with the negative component in the direction normal to $E_{i,N}$. We also have an extension

$$(4.2) \quad 0 \rightarrow \mathcal{N}_{\ell_j/W} \rightarrow \mathcal{N}_h|_{\ell_j} \rightarrow Q(\ell_j) \rightarrow 0,$$

where $Q(\ell_j)$ is a torsion sheaf supported at the points where ℓ_j meets the adjacent components. For $j = N$ these are $g'_i(\mathbb{P}^1)$ and ℓ_{N-1} , and since the tangent vectors to these curves are normal to $E_{i,N}$, we find

$$\mathcal{N}_h|_{\ell_N} \simeq \mathcal{O}_{\mathbb{P}^1}(1)^{\dim(V)-2} \oplus \mathcal{O}_{\mathbb{P}^1}(1).$$

The normal bundle of the proper transform of a line in the blow-up of projective space at a point of the line is

$$\mathcal{N}_{\ell_j/E_{i,j}} = \mathcal{N}_{\ell_j/\text{Bl}_{w_{i,j}}\mathbb{P}^{\dim(V)-1}} \simeq \mathcal{O}_{\mathbb{P}^1}^{\dim(V)-2}$$

for $j = 1, \dots, N-1$. Similarly, we can compute

$$\mathcal{N}_{E_{i,h}/W}|_{\ell_j} = \mathcal{O}_{\mathbb{P}^1}(-2)$$

so the exact sequence analogous to (4.1) yields

$$\mathcal{N}_{\ell_j/W} \simeq \mathcal{O}_{\mathbb{P}^1}^{\dim(V)-2} \oplus \mathcal{O}_{\mathbb{P}^1}(-2),$$

with the negative component in the direction normal to $E_{i,j}$. Using (4.2) and the fact that ℓ_j is adjacent to ℓ_{j+1} and ℓ_{j-1} (or $g''_i(\mathbb{P}^1)$ when $j = 1$), we find

$$\mathcal{N}_h|_{\ell_j} \simeq \mathcal{O}_{\mathbb{P}^1}^{\dim(V)-2} \oplus \mathcal{O}_{\mathbb{P}^1}.$$

□

Definition 14. A smooth separably rationally connected variety Y is *strongly rationally connected* if any of the following conditions hold:

- (1) for each point $y \in Y$, there exists a rational curve $f : \mathbb{P}^1 \rightarrow Y$ joining y and a generic point in Y ;
- (2) for each point $y \in Y$, there exists a free rational curve containing y ;

- (3) for any finite collection of points $y_1, \dots, y_m \in Y$, there exists a very free rational curve containing the y_j as smooth points;
- (4) for any finite collection of jets

$$\text{Spec}(k[\epsilon]/\langle \epsilon^{N+1} \rangle) \subset Y, i = 1, \dots, m$$

supported at distinct points y_1, \dots, y_m , there exists a very free rational curve smooth at y_1, \dots, y_m and containing the prescribed jets.

The implications

$$(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow (1)$$

are obvious. By Proposition 10, assertions (1)-(3) are each equivalent to the condition $Y = Y^0$. Property (4) is analogous to Theorem 2.2 of [7], which is stated for proper varieties. It follows from (1) by Proposition 13.

With basic properties of strongly rationally connected varieties established, Theorem 1 follows from the general result (cf. [15] IV.6.10.1):

Theorem 15. *Let $\pi : \mathcal{Y} \rightarrow B$ be a smooth morphism whose fibers are strongly rationally connected. Assume that π has a section. Then sections of $\mathcal{Y} \rightarrow B$ satisfy approximation away from S .*

Proof. Let $\bar{\pi} : \bar{\mathcal{Y}} \rightarrow \bar{B}$ be a proper model of $\mathcal{Y} \rightarrow B$, which exists by [20]. The section extends to a section \bar{s} of $\bar{\pi}$. By a result of Artin and Néron [1] Corollary 4.6, there exists a blow-up with center supported in $\bar{\pi}^{-1}(S)$

$$\tilde{\mathcal{Y}} \rightarrow \bar{\mathcal{Y}}$$

such that the proper transform of $\bar{s}(\bar{B})$ in $\tilde{\mathcal{Y}}$ is contained in $\tilde{\mathcal{Y}}^{sm}$.

Recall the proof of weak approximation at places of good reduction in Section 5 of [10]. This is a bootstrap argument, using the existence of a section in the smooth locus to construct sections with prescribed jets of successively higher order. For the base case, suppose we are given an arbitrary section $t : B \rightarrow \mathcal{Y}$. When $\mathcal{Y} \rightarrow B$ is proper, Kollar-Miyaoka-Mori [16] first demonstrated how to get a section with prescribed values y_1, \dots, y_r at $b_1, \dots, b_r \in B$. The key is to construct a comb with handle $t(B)$ and teeth very free curves in the fibers \mathcal{Y}_{b_i} joining $t(b_i)$ to y_i , which deforms to a section passing through the y_i . For the inductive step, suppose we have a section with prescribed jets to order $\leq N-1$ at a finite set of points b_1, \dots, b_r . Blow up the total space N times along the jet data at each of b_1, \dots, b_r to get a new model with (admittedly very special) *reducible* fibers over b_1, \dots, b_r . The N th-order jet conditions

in the original model translate into point conditions in the new model. There we produce an explicit comb with reducible teeth, based on the proper transform of the section obtained by the inductive hypothesis, that deforms to the desired section. This only requires the existence of very free curves in \mathcal{Y}_{b_i} passing through y_i with prescribed tangency.

Properness is used twice. At the zeroth-order step, it is used to exhibit the very free fibral curves joining $t(b_i)$ to y_i . In the inductive step, it is used to find a very free fibral curve with prescribed tangency at y_i . In our situation, these are guaranteed by the hypothesis that the fibers are strongly rationally connected. \square

5. CUBIC SURFACES

We work over an algebraically closed field of characteristic zero.

Definition 16. A *log Del Pezzo surface* is a pair (X, Δ) consisting of a normal projective surface X and an effective \mathbb{Q} -divisor $\Delta = \sum a_i \Delta_i$, $0 < a_i \leq 1$ on X , with log terminal singularities, such that $-(K_X + \Delta)$ is ample. When Δ is empty, this is equivalent to saying that X has quotient singularities and ample anticanonical class.

Theorem 17 ([11] 1.6). *The smooth locus of a log Del Pezzo surface (X, Δ) is rationally connected, i.e., two generic points in X^{sm} can be joined by an irreducible projective rational curve contained in X^{sm} .*

Example 18 ([23]). There exist projective rational surfaces with rational double points whose smooth locus is not rationally connected. Consider

$$\tilde{X} = E \times \mathbb{P}^1$$

where $(E, 0)$ is an elliptic curve and the involution

$$\begin{aligned} \iota : \tilde{X} &\rightarrow \tilde{X} \\ (e, [x_0, x_1]) &\mapsto (-e, [x_1, x_0]). \end{aligned}$$

The involution has eight isolated fixed points $\mathfrak{q} \subset \tilde{X}$. The quotient $X = \tilde{X}/\langle \iota \rangle$ has eight A_1 singularities and is rational: $X \rightarrow E/\langle \iota \rangle \simeq \mathbb{P}^1$ is a conic bundle. Since $\tilde{X} - \mathfrak{q} \rightarrow X^{sm}$ is a covering space, $\pi_1(X^{sm}) \subset \pi_1(\tilde{X} - \mathfrak{q})$ with index two. Thus

$$\pi(\tilde{X} - \mathfrak{q}) \simeq \pi(\tilde{X}) \simeq \pi(E) \simeq \mathbb{Z} \times \mathbb{Z}$$

and X^{sm} has infinite fundamental group. However, rationally connected varieties (even non-proper ones) have finite fundamental groups (see Lemma 7.8 of [11] and Proposition 2.10 of [14], for example).

The following conjecture would allow us to apply Theorem 1 to prove weak approximation for many log Del Pezzo surfaces:

Conjecture 19. The smooth locus of a log Del Pezzo surface is strongly rationally connected.

We prove this for cubic surfaces:

Theorem 20. *Let $X \subset \mathbb{P}^3$ be a cubic surface with rational double points. Then X^{sm} is strongly rationally connected.*

Proof. Let $x_1 \in X^{sm}$ be a point. We produce a rational curve $R \subset X^{sm}$ joining x_1 and a generic point $x_2 \in X^{sm}$.

We start with an elementary lemma:

Lemma 21. *Let $Y \subset \mathbb{P}^n$ be an irreducible hypersurface such that the Gauss map*

$$\begin{aligned} Y &\dashrightarrow \check{\mathbb{P}}^n \\ y &\mapsto [\mathcal{T}_Y|_y] \end{aligned}$$

is generically finite. Then a generic tangent hyperplane section to Y has an isolated singularity of multiplicity two with smooth projectivized tangent cone.

Proof. Since the Gauss map is generically finite, its differential is generically of maximal rank. However, the differential at $y \in Y$ can be identified with the dual to the second fundamental form (see [9, 17.11])

$$\Pi_y : \text{Sym}^2(\mathcal{T}_Y|_y) \rightarrow \mathcal{N}_{Y/\mathbb{P}^n}|_y.$$

This is nondegenerate precisely when the quadratic term of the Taylor expansion of the defining equation of the tangent hyperplane section H_y has maximal rank. \square

This is applicable to cubic surfaces X with rational double points. It is a classical fact that X contains a finite number of lines. However, if the image of the Gauss map of X is a curve C then X is dual to C and thus ruled by lines.

Now we will make explicit how x_2 must be chosen. Applying the lemma, we may assume

(1) The tangent hyperplane section H_2 at x_2 is irreducible and nodal.

In particular, $H_2 \subset X^{sm}$ and there are no lines $\ell \subset X$ containing x_2 .

Projection from x_2 then gives a double cover

$$\text{Bl}_{x_2} X \rightarrow \mathbb{P}^2;$$

the covering transformation interchanges the exceptional divisor and the proper transform. We obtain a birational involution

$$\begin{aligned} \iota_{x_2} : X &\dashrightarrow X \\ x &\mapsto x', \end{aligned}$$

where $\{x, x', x_2\}$ are collinear. This factors as the blow-up of x_2 followed by the blow-down of the proper transform of H_2 . Note that ι_{x_2} fixes the singularities of X and thus takes X^{sm} to itself.

We also assume:

(2) H_2 does not contain x_1 .

It follows that H_2 does not contain $x'_1 = \iota_{x_2}(x_1)$. Moreover, x_1 and x'_1 are in the open subset on which ι_{x_2} is an isomorphism.

We assume furthermore:

(3) x_2 is not contained in H_1 , the tangent hyperplane to X at x_1 .

It follows that $x_2 \notin H'_1$, the tangent hyperplane section at x'_1 . Indeed, suppose that $x_2 \in H'_1$. We know that $x_2 \neq x'_1$ (because $x'_1 \notin H_2$), so consider the line joining x_2 and x'_1 . This meets X only at x_2 and x'_1 , so $x'_1 = x_1$ and $x_2 \in H_1$, a contradiction.

Finally, we assume:

(4) H'_1 is irreducible and nodal.

In particular, $H'_1 \subset X^{sm}$.

Since $x_2 \notin H'_1$, ι_{x_2} is regular along H'_1 . We verify that the rational curve $R = \iota_{x_2}(H'_1)$ has the desired properties. Since $x_2 \notin H'_1$, H_2 and H'_1 intersect at a point $z \neq x_2$; thus the curve

$$R = \iota_{x_2}(H'_1) \ni \iota_{x_2}(z) = x_2.$$

We know $H'_1 \subset X^{sm}$ and $\iota_{x_2}(X^{sm}) \subset X^{sm}$, hence $R \subset X^{sm}$. We have $x'_1 \in H'_1$, so $x_1 = \iota_{x_2}(x'_1) \in R$. Since H'_1 meets H_2 in a point $z \neq x_2$, $x_2 = \iota_{x_2}(y) \in R$. \square

We now prove Theorem 2: For each singular fiber \mathcal{X}_b , \mathcal{X}_b^{sm} is strongly rationally connected by Theorem 20. Approximation follows from Theorem 1.

Example 22. Here is another case where Conjecture 19 is easily verified. Let X be a partial resolution of a cubic surface Σ with at most A_1 -singularities, i.e., we have a factorization of the minimal resolution

$$\tilde{\Sigma} \rightarrow X \xrightarrow{\beta} \Sigma.$$

Then X^{sm} is strongly rationally connected.

Theorem 2 implies that Σ^{sm} is strongly rationally connected, hence $\beta^{-1}(\Sigma^{sm}) \subset (X^{sm})^0$. The locus $X^{sm} \setminus \beta^{-1}(\Sigma^{sm})$ is a union of (-2) -curves $\{E_i\}$, corresponding to the resolved singularities $\{p_i\}$ of Σ . If $(X^{sm})^0$ meets E_i , it must also contain E_i . Hence it suffices to show that for each E_i there exists a rational curve in X^{sm} meeting E_i and $\beta^{-1}(\Sigma^{sm})$ (see Remark 11).

To find this rational curve, consider the projection from p_i

$$\pi_i : \Sigma \dashrightarrow \mathbb{P}^2$$

which induces a morphism $\pi'_i : X \rightarrow \mathbb{P}^2$. The image of E_i is a plane conic and the image of the singularities of X has codimension two in \mathbb{P}^2 , so there exists a rational curve

$$f : \mathbb{P}^1 \rightarrow \mathbb{P}^2 \setminus \pi'_i(\text{Sing}(X))$$

meeting the image of E_i .

The same argument applies if X is obtained from a cubic surface Σ with A_1 and A_2 singularities by resolving some subset of $\text{Sing}(\Sigma)$.

Corollary 4 is an immediate consequence of Corollary 3 and the following:

Lemma 23. *Let $\mathcal{Hilb} = \mathbb{P}(\Gamma(\mathcal{O}_{\mathbb{P}^n}(d)))$ denote the Hilbert scheme of degree- d hypersurfaces, $\mathcal{U} \rightarrow \mathcal{Hilb}$ the universal family, and $D \subset \mathcal{Hilb}$ the discriminant divisor. Suppose $B \subset \mathcal{Hilb}$ is a smooth curve with corresponding family*

$$\mathcal{Y} := \mathcal{U} \times_{\mathcal{Hilb}} B \rightarrow B.$$

Then B is transversal to each branch of the discriminant iff \mathcal{Y} is regular and the fibers have ordinary double points.

Our transversality condition means that each branch of D at b is smooth and transverse to B .

Proof. Let $b \in B \cap D$ and $y \in Y_b$ a singularity corresponding to a branch $D' \subset D$ at b . First suppose that y is an isolated singularity of \mathcal{Y}_b . Then we have the formula [22, 2.8.3]

$$\text{mult}_b(B \cap D') = \mu(\mathcal{Y}_b, y) + \mu(\mathcal{Y}, y),$$

the sum of the corresponding Milnor numbers. Thus the multiplicity is one if $\mu(\mathcal{Y}_b, y) = 1$ and $\mu(\mathcal{Y}, y) = 0$, i.e., $y \in \mathcal{Y}$ is nonsingular and $y \in \mathcal{Y}_b$ is an ordinary double point.

Now suppose y fails to be isolated. If $d = 2$ then the quadratic form defining \mathcal{Y}_b has rank $\leq n - 1$; the discriminant is defined by the determinant of an $(n + 1) \times (n + 1)$ symmetric matrix and thus is singular when the matrix has rank $< n$. Otherwise, let $\Sigma \subset \mathcal{Y}_b$ denote the irreducible component of the singular locus containing y . Choose generic $y', y'' \in \Sigma, y' \neq y''$, and consider the hypersurfaces singular at both y' and y'' , which form a codimension-2($n + 1$) linear subspace $L \subset \mathcal{H}ilb = \mathbb{P}(\Gamma(\mathcal{O}_{\mathbb{P}^n}(d)))$. The generic hypersurface singular at y' (or y'') is contained in D' thus L is contained in the singular locus of D' . \square

6. HIGHER-DIMENSIONAL FANO HYPERSURFACES

Here we work over an uncountable algebraically closed field k of characteristic zero.

We are grateful to James McKernan for pointing out the following amplification of [11, 5.9]

Proposition 24. *Let X be a projective rationally connected variety with isolated terminal local complete intersection singularities. Then X^{sm} is strongly rationally connected.*

Proof. Let $\rho : \tilde{X} \rightarrow X$ denote a resolution of singularities of X , such that $\rho^{-1}(X^{sm}) \rightarrow X^{sm}$ is an isomorphism and $\tilde{X} \setminus \rho^{-1}(X^{sm})$ is a normal crossings divisor with components E_1, \dots, E_m .

We first show that X^{sm} is rationally connected. Suppose that $(x_1, x_2) \in X^{sm} \times X^{sm}$ is general, in the sense that it lies in the complement of a countable union of proper subvarieties. (Here we are using the fact that the base field is uncountable.) Then *any* morphism

$$h : \mathbb{P}^1 \rightarrow \tilde{X}, h(0) = x_1, h(\infty) = x_2$$

is necessarily very free (cf. [15, 3.11]).

Choose a very free imbedding $\tilde{f} : \mathbb{P}^1 \rightarrow \tilde{X}$ with $\tilde{f}(0) = x_1, \tilde{f}(\infty) = x_2$, and image meeting $\rho^{-1}(X^{sm})$. The induced curve in X is denoted $f = \rho \circ \tilde{f} : \mathbb{P}^1 \rightarrow X$. We may assume that $f(\mathbb{P}^1)$ meets the singularities of X ; otherwise there is nothing to prove. Consequently, $\tilde{f}(\mathbb{P}^1)$ meets at least one of the E_i .

We compare dimensions of deformation spaces

$$\text{Def}(\tilde{f}) := \text{Hom}(\mathbb{P}^1, \tilde{X}; 0 \rightarrow x_1, \infty \rightarrow x_2)$$

and

$$\text{Def}(f) := \text{Hom}(\mathbb{P}^1, X; 0 \rightarrow x_1, \infty \rightarrow x_2).$$

The first space has dimension

$$-\deg \tilde{f}^*K_{\tilde{X}} - 2\dim(X)$$

at \tilde{f} . By Theorem 2.10 of [13], the second space has dimension at least

$$-\deg f^*K_X - 2\dim(X);$$

the discrepancy formula

$$K_{\tilde{X}} = K_X + \sum_i b_i E_i, \quad b_i > 0$$

then guarantees

$$\dim \text{Def}(\tilde{f}) < \dim \text{Def}(f).$$

Composition by ρ gives a morphism of deformation spaces

$$\begin{aligned} \iota_\rho : \text{Def}(\tilde{f}) &\rightarrow \text{Def}(f) \\ \tilde{g} &\mapsto \rho \circ \tilde{g} \end{aligned}$$

which is not dominant by dimension considerations. Let $g_t : \mathbb{P}^1 \rightarrow X$ be a one-parameter deformation of f such that

$$\lim_{t \rightarrow 0} g_t = f,$$

and $g_t(0) = x_1, g_t(\infty) = x_2$ for each t . Assume that g_t is generic on the maximal-dimension irreducible component of $\text{Def}(f)$ passing through f . For generic t , write

$$\tilde{g}_t : \mathbb{P}^1 \rightarrow \tilde{X}$$

is the lift of g_t to \tilde{X} , which is also very free.

We claim $g_t(\mathbb{P}^1) \subset X^{sm}$. If not then we could repeat the argument above, finding a component of $\text{Def}(g_t)$ with dimension strictly larger than the dimension of $\text{Def}(\tilde{g}_t)$, contradicting our assumption.

We now prove that X^{sm} is *strongly* rationally connected, by exhibiting a free curve in X^{sm} through each point $x_1 \in X^{sm}$. As above, let $\tilde{f} : \mathbb{P}^1 \rightarrow \tilde{X}$ denote a free curve with $\tilde{f}(0) = x_1$ and passing through a general point of \tilde{X} . Write $f = \rho \circ \tilde{f}$ and repeat our dimension analysis, applied to the deformation spaces

$$\text{Def}(\tilde{f}) := \text{Hom}(\mathbb{P}^1, \tilde{X}; 0 \rightarrow x_1)$$

and

$$\text{Def}(f) := \text{Hom}(\mathbb{P}^1, X; 0 \rightarrow x_1).$$

If $f(\mathbb{P}^1)$ is not contained in X^{sm} then

$$\dim \text{Def}(\tilde{f}) < \dim \text{Def}(f)$$

and we can choose $g_t : \mathbb{P}^1 \rightarrow X$ generic on the component of maximal dimension. The lift $\tilde{g}_t : \mathbb{P}^1 \rightarrow \tilde{X}$ remains free because it passes through a general point of \tilde{X} . Thus if $g_t(\mathbb{P}^1)$ were not in X^{sm} then we could exhibit a component of $\text{Def}(g_t)$ with dimension strictly larger than the dimension of $\text{Def}(\tilde{g}_t)$, a contradiction. \square

Remark 25. The classification of terminal singularities in dimension three [19] shows they are quotients of isolated complete intersection singularities by the action of $\mathbb{Z}/r\mathbb{Z}$, where r is the index of the singularity. Thus the hypotheses of Proposition 24 are quite natural.

With further technical hypotheses, the proof of Proposition 24 can be extended to $r > 1$.

Example 26. Examples of three-dimensional terminal singularities of index one include ordinary threefold double points

$$w^2 = x^2 + y^2 + z^2.$$

For a complete list see [17, 6.4].

Theorem 1 then gives

Corollary 27. *Let X be a smooth rationally connected variety over $F = k(B)$. Suppose X admits a regular proper model $\pi : \mathcal{X} \rightarrow B$ whose singular fibers have isolated terminal complete-intersection singularities. Then weak approximation holds for X away from $S = \overline{B} \setminus B$.*

Applying Lemma 23 we obtain

Corollary 28. *Let $\mathcal{Hilb} = \mathbb{P}(\Gamma(\mathcal{O}_{\mathbb{P}^n}(d))) \simeq \mathbb{P}^{\binom{n+d}{d}-1}$ denote the Hilbert scheme of hypersurfaces of degree $d \leq n, n \geq 4$, $\mathcal{U} \rightarrow \mathcal{Hilb}$ the universal family, and $D \subset \mathcal{Hilb}$ the discriminant divisor. Let $B \subset \mathcal{Hilb}$ be a smooth curve transverse to each branch of D and*

$$\mathcal{X} := \mathcal{U} \times_{\mathcal{Hilb}} B \rightarrow B$$

the corresponding family. Then $X = \mathcal{U} \times_{\mathcal{Hilb}} \text{Spec}(F)$ satisfies weak approximation away from $S = \overline{B} \setminus B$.

REFERENCES

- [1] M. Artin. Algebraic approximation of structures over complete local rings. *Inst. Hautes Études Sci. Publ. Math.*, (36):23–58, 1969.
- [2] Fedor Bogomolov and Yuri Tschinkel. Rational curves and points on $K3$ surfaces. *Amer. J. Math.*, 127(4):825–835, 2005.

- [3] E. Brieskorn. Die Auflösung der rationalen Singularitäten holomorpher Abbildungen. *Math. Ann.*, 178:255–270, 1968.
- [4] E. Brieskorn. Singular elements of semi-simple algebraic groups. In *Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 2*, pages 279–284. Gauthier-Villars, Paris, 1971.
- [5] J. L. Colliot-Thélène and P. Gille. Remarques sur l’approximation faible sur un corps de fonctions d’une variable. In *Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002)*, volume 226 of *Progr. Math.*, pages 121–134. Birkhäuser Boston, Boston, MA, 2004.
- [6] A. J. de Jong and J. Starr. Low degree complete intersections are rationally simply connected, 2006. preprint.
- [7] O. Debarre. Variétés rationnellement connexes (d’après T. Graber, J. Harris, J. Starr et A. J. de Jong). *Astérisque*, 290:exposé 905, ix, 243–266, 2003. Séminaire Bourbaki. Vol. 2001/2002.
- [8] T. Graber, J. Harris, and J. Starr. Families of rationally connected varieties. *J. Amer. Math. Soc.*, 16(1):57–67, 2003.
- [9] J. Harris. *Algebraic geometry*, volume 133 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [10] B. Hassett and Y. Tschinkel. Weak approximation over function fields. *Inventiones mathematicae*, 163:171–190, 2006.
- [11] S. Keel and J. McKernan. Rational curves on quasi-projective surfaces. *Mem. Amer. Math. Soc.*, 140(669):viii+153, 1999.
- [12] A. Knecht. *Approximation at Places of Bad Reduction for Degree Two del Pezzo Surfaces*. PhD thesis, Rice University, Houston, Texas, 2007.
- [13] J. Kollar. Cone theorems and bug-eyed covers. *J. Algebraic Geom.*, 1(2):293–323, 1992.
- [14] J. Kollar. *Shafarevich maps and automorphic forms*. M. B. Porter Lectures. Princeton University Press, Princeton, NJ, 1995.
- [15] J. Kollar. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Math.* Springer-Verlag, Berlin, 1996.
- [16] J. Kollar, Y. Miyaoka, and S. Mori. Rationally connected varieties. *J. Algebraic Geom.*, 1(3):429–448, 1992.
- [17] J. Kollar and N. I. Shepherd-Barron. Threefolds and deformations of surface singularities. *Invent. Math.*, 91(2):299–338, 1988.
- [18] D. Madore. Approximation faible aux places de bonne réduction sur les surfaces cubiques sur les corps des fonctions, 2004. preprint, to appear.
- [19] S. Mori. On 3-dimensional terminal singularities. *Nagoya Math. J.*, 98:43–66, 1985.
- [20] M. Nagata. A generalization of the imbedding problem of an abstract variety in a complete variety. *J. Math. Kyoto Univ.*, 3:89–102, 1963.
- [21] P. Swinnerton-Dyer. Weak approximation and R -equivalence on cubic surfaces. In *Rational points on algebraic varieties*, volume 199 of *Progr. Math.*, pages 357–404. Birkhäuser, Basel, 2001.

- [22] B. Teissier. Introduction to equisingularity problems. In *Algebraic geometry (Proc. Sympos. Pure Math., Vol. 29, Humboldt State Univ., Arcata, Calif., 1974)*, pages 593–632. Amer. Math. Soc., Providence, R.I., 1975.
- [23] D.-Q. Zhang. Algebraic surfaces with nef and big anti-canonical divisor. *Math. Proc. Cambridge Philos. Soc.*, 117(1):161–163, 1995.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, MS 136, HOUSTON, TEXAS 77251-1892, USA

E-mail address: hassett@rice.edu

MATHEMATISCHES INSTITUT, BUNSENSTR. 3-5, 37073 GÖTTINGEN, GERMANY,
COURANT INSTITUTE, NYU, 251 MERCER STR., NEW YORK, NY 10012, USA

E-mail address: tschinkel@cims.nyu.edu