

MODELS OF BRAUER-SEVERI SURFACE BUNDLES

ANDREW KRESCH AND YURI TSCHINKEL

1. INTRODUCTION

This paper is motivated by the study of rationality properties of Mori fiber spaces. These are algebraic varieties, naturally occurring in the minimal model program, typically via contractions along extremal rays; they are fibrations with *geometrically* rational generic fiber. Widely studied are conic bundles $\pi : X \rightarrow S$, when the generic fiber is a conic. According to the Sarkisov program [70], there exists a birational modification

$$\begin{array}{ccc} \tilde{X} & \dashrightarrow & X \\ \tilde{\pi} \downarrow & & \downarrow \pi \\ \tilde{S} & \longrightarrow & S \end{array}$$

such that:

- (i) the general fiber of $\tilde{\pi}$ is a smooth conic,
- (ii) the discriminant divisor of $\tilde{\pi}$ is a simple normal crossing divisor,
- (iii) generally along the discriminant divisor, the fiber is a union of two lines, and
- (iv) over the singular locus of the discriminant divisor, the fiber is a double line in the plane.

Rationality of conic bundles over surfaces has been classically studied by Artin and Mumford [12], who produced examples of nonrational unirational threefolds of this type, computing their Brauer groups and using its nontriviality as an obstruction to rationality. This was generalized to higher-dimensional quadric bundles in [22], bringing higher unramified cohomology into the subject and providing further examples of nonrational varieties based on these new obstructions.

The specialization method, introduced by Voisin [75] and developed further by Colliot-Thélène–Pirutka [23], emerged as a powerful tool in the study of *stable rationality*. It allows to relate the failure of stable rationality of a very general member of a flat family to the existence of special fibers with nontrivial unramified cohomology and mild singularities; see also [15], [74]. In particular, the stable rationality problem for

Date: December 16, 2018.

very general smooth rationally connected threefolds can be reduced to the case of conic bundles over rational surfaces [39], [49]. The case of very general families of conic bundles over rational surfaces is treated in [36]. Many families that arise naturally in applications lead to discriminant curves of a special form. These have been studied, e.g., in [18] and [39].

A key step in the specialization method is the construction of families of Mori fiber spaces with controlled degeneration. A next case to consider is a fibration whose generic fiber is a form of \mathbb{P}^2 .

Definition 1.1. Let k be a perfect field of characteristic different from 3 and $\pi: X \rightarrow S$ a morphism of smooth proper varieties over k . We call π a *standard Brauer-Severi surface bundle* if there exists a reduced divisor $D \subset S$ whose singular locus D^{sing} is smooth, such that:

- (i) π is smooth over $S \setminus D$ and the generic fiber of π is a nontrivial form of \mathbb{P}^2 over $k(S)$.
- (ii) Over every geometric point of $D \setminus D^{\text{sing}}$ the fiber of π is a union of three Hirzebruch surfaces \mathbb{F}_1 , meeting transversally, such that any pair of them meets along a fiber of one and the (-1) -curve of the other, while over the generic point of every irreducible component of D the fiber of π is irreducible.
- (iii) Over every geometric point of D^{sing} the fiber of π is an irreducible scheme whose underlying reduced subscheme is isomorphic to the cone over a twisted cubic curve.

Since a morphism of smooth varieties with fibers all of the same dimension is flat [57, Thm. 23.1], $\pi: X \rightarrow S$ as in Definition 1.1 is necessarily flat. There is a geometric description, which in the case of algebraically closed base field is due to Artin [9], [10], and Maeda [55] (which we state with a less restrictive hypothesis on the base field):

Theorem 1.2. *Let k be a perfect field of characteristic different from 3 and S a smooth proper algebraic variety over k . Assume that over k we have resolution of marked ideals on smooth varieties of dimension equal to $\dim(S)$. Let*

$$\pi: X \rightarrow S$$

be a morphism of proper varieties over k , whose generic fiber is a nontrivial form of \mathbb{P}^2 . Then there exists a commutative diagram

$$\begin{array}{ccc} \widetilde{X} & \xrightarrow{\varrho_X} & X \\ \tilde{\pi} \downarrow & & \downarrow \pi \\ \widetilde{S} & \xrightarrow{\varrho_S} & S \end{array}$$

where, if we let $U \subset S$ denote the locus over which π is smooth,

- ϱ_S is a birational morphism that restricts to an isomorphism over U ,
- ϱ_X is a birational map that restricts to an isomorphism over $\pi^{-1}(U)$, and
- $\tilde{\pi}$ is a standard Brauer-Severi surface bundle with singular fibers over a simple normal crossing divisor in \tilde{S} .

We note that a standard Brauer-Severi surface bundle with singular fibers over a simple normal crossing divisor $D = D_1 \cup \dots \cup D_\ell$ necessarily has $D_i \cap D_{i'} \cap D_{i''} = \emptyset$ for any triple of distinct indices i, i', i'' . Statements on conic bundles, analogous to Theorem 1.2, have appeared in [70] and more recently in [64].

Remark 1.3. When $\text{char}(k) = 0$ or $\dim(S) = 2$, the resolution hypothesis stated in Theorem 1.2 is standard. For the notion of resolution of marked ideals, the reader may consult [76]; this procedure ensures that any boundary may be converted to a simple normal crossing divisor by a sequence of blow-ups along smooth centers, which may furthermore be chosen to have normal crossing with a given simple normal crossing divisor on S .

In this paper we describe a technique based on root stacks, appearing in [36], that allows us to recover the Sarkisov program and its version by Artin and Maeda (Theorem 1.2) and obtain an extension [48] to more general del Pezzo surface fibrations; a related work is [50]. The proof of Theorem 1.2 illustrates this technique, which leads to a global version of the constructions of Artin and Maeda that is crucial in moduli problems and in applications of specialization. As an application to rationality problems we prove:

Theorem 1.4. *Let S be a smooth del Pezzo surface of degree 2 over an uncountable algebraically closed field k of characteristic $\neq 2, 3$, and let $d \geq 3$. Then a standard Brauer-Severi surface bundle ramified over a very general member of the linear system $|d(-K_S)|$ is not stably rational.*

Remark 1.5. Properties of Brauer groups, combined with Theorem 1.2, ensure the existence of Brauer-Severi surface bundles as in Theorem 1.4. Details are presented in Section 10.

More sophisticated methods [47] extend Theorem 1.4 to $d \geq 2$; the threshold $|2K_S + D| \neq \emptyset$ ($D \in |d(-K_S)|$), coincides with that in Shokurov's conjecture on rationality of conic bundles [71]. Even the non-rationality of the fourfolds considered in Theorem 1.4 was unknown; indeed, the standard obstructions to rationality of threefolds cannot be adapted to treat these fourfolds; cf. [14]

Much recent progress in the study of rationality relies on degenerations to reference varieties which, in the case of threefolds, are conic bundles over rational surfaces [39] [37]. In the study of rationality properties of fourfolds, quadric surface bundles [38] and more general del Pezzo surface bundles [3] played an important role; this motivates our investigation of Brauer-Severi surface bundles.

Acknowledgments: We are grateful to the referee for many helpful suggestions, to Brendan Hassett and Alena Pirutka for stimulating discussions on related topics, to Asher Auel for helpful comments, and to David Rydh for the streamlined construction of the birational modification in Section 9. The first author is partially supported by the Swiss National Science Foundation. The second author is partially supported by NSF grant 1601912.

2. PRELIMINARIES

We start with some algebraic results that will be used in our approach. Let k be a field. By a *variety* over k we mean a geometrically integral separated finite-type scheme over k . We will need, however, the generality of locally Noetherian schemes and Deligne-Mumford stacks. This additional generality allows us to construct models of Brauer-Severi bundles in which all of the fibers are smooth. This comes at the cost of imposing nontrivial stack structure on the base.

All cohomology in this paper is étale cohomology.

2.1. Deligne-Mumford stacks. Deligne and Mumford [25] defined a class of stacks which includes all stack quotients of the form $[X/G]$, where X is a variety and G a finite group, and which are now called Deligne-Mumford stacks. The stack $[X/G]$ differs from the conventional quotient X/G (which exists as a variety, e.g., when X is quasi-projective) in that it keeps track of the stabilizers of the G -action. A further advantage is that $[X/G]$ has the same local properties (smoothness, etc.) as X . This will be crucial for us, as we will rely on properties of smooth varieties. Most important for us will be *orbifolds* over k , which are smooth geometrically integral separated finite-type Deligne-Mumford stacks over k that possess an open substack isomorphic to a scheme.

As a stack, a scheme X is encoded by the category of all schemes with morphism to X , and $[X/G]$, as G -torsors $E \rightarrow T$ with equivariant morphism $E \rightarrow X$. When X is a point (i.e., $X = \text{Spec}(k)$ if we work over a base field k , or $\text{Spec}(\mathbb{Z})$ if we work with all schemes), $[X/G]$ is the

category BG of G -torsors, the *classifying stack* of G . We observe, the diagonal of BG is a finite étale morphism of degree $|G|$.

An infinite discrete group G is also permitted according to modern usage of the term Deligne-Mumford stack, leading to pathologies such as zero-dimensional Deligne-Mumford stack $[\mathrm{Spec}(\overline{\mathbb{Q}} \otimes_{\mathbb{Q}} \overline{\mathbb{Q}})/H]$ for an index 2 non-open subgroup H of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, which exists (by [72, Thm. 1]), is reduced (by [32, 4.3.5]), and is irreducible with 2 points (objects over $\mathrm{Spec}(\Omega)$ for all fields Ω , modulo morphisms over $\mathrm{Spec}(\Omega') \rightarrow \mathrm{Spec}(\Omega)$ for embeddings $\Omega \rightarrow \Omega'$).

When we restrict to Noetherian Deligne-Mumford stacks, i.e., quasi-compact locally Noetherian Deligne-Mumford stacks with quasi-compact diagonal, such pathologies are excluded.

More general algebraic stacks such as BG for positive-dimensional algebraic groups or nonreduced group schemes G (Artin stacks) are not needed here.

2.2. Gerbes. Let X be a Noetherian Deligne-Mumford stack and n a positive integer, invertible in the local rings of an étale atlas of X . A gerbe over X banded by roots of unity μ_n , or just μ_n -gerbe, is a Deligne-Mumford stack \mathcal{G} with morphism $\mathcal{G} \rightarrow X$ that étale locally over X is isomorphic to a product with the classifying stack $B\mu_n$ and is equipped with compatible identifications of the automorphism groups of local sections with μ_n .

A μ_n -gerbe is classified, up to isomorphism compatible with the identifications of automorphism groups of local sections, by a class in $H^2(X, \mu_n)$; cf. [59, §IV.2].

As mentioned in [45, §3], a vector bundle on a gerbe banded by roots of unity with scalar action by stabilizers determines a Brauer-Severi fibration. Specifically, the projectivization $\mathbb{P}(E)$ of a given vector bundle descends via $\mathcal{G} \rightarrow X$ to yield smooth projective $P \rightarrow X$ with $\mathcal{G} \times_X P \cong \mathbb{P}(E)$. This framework was essential in [36], in the construction of families of conic bundles for the application of the specialization method. We will use the same strategy for families of Brauer-Severi bundles of relative dimension 2.

2.3. Brauer groups. The Brauer group $\mathrm{Br}(K)$ of a field K is a classical invariant, defined as the group of similarity classes of central simple algebras over K . It may also be described in terms of Galois cohomology, or in the language of étale cohomology, as

$$\mathrm{Br}(K) \cong H^2(\mathrm{Spec}(K), \mathbb{G}_m).$$

Now let S be a Noetherian scheme or Deligne-Mumford stack. Similarity classes of sheaves of Azumaya algebras over S , which naturally

generalize central simple algebras over a field, give one notion of Brauer group of S . Another, more relevant for us, is the *cohomological Brauer group*

$$\mathrm{Br}(S) := H^2(S, \mathbb{G}_m)_{\mathrm{tors}}.$$

The former is a subgroup of the latter; despite examples of a pathological nature where they differ [27] they are known to coincide in geometrically relevant cases, including all schemes that possess an ample line bundle; see, e.g., [53] and references therein.

Proposition 2.1 ([33, Cor. II.1.8, II.2.2], [36, Prop. 2]). *Let S be a regular integral Noetherian scheme or a regular integral Noetherian Deligne-Mumford stack, and $S' \subset S$ a nonempty open subscheme, respectively, substack. Then:*

- (i) *The group $H^2(S, \mathbb{G}_m)$ is torsion, and the restriction homomorphism $\mathrm{Br}(S) \rightarrow \mathrm{Br}(S')$ is injective.*
- (ii) *If $\dim(S) \leq 2$ then every element of $\mathrm{Br}(S)$ is the class of a sheaf of Azumaya algebras on S , and every sheaf of Azumaya algebras on S' representing an element $\alpha|_{S'}$ for some $\alpha \in \mathrm{Br}(S)$ extends to a sheaf of Azumaya algebras on S .*
- (iii) *If $\dim(S) = 2$ and $\dim(S \setminus S') = 0$ then every sheaf of Azumaya algebras on S' extends to a sheaf of Azumaya algebras on S , and the restriction homomorphism $\mathrm{Br}(S) \rightarrow \mathrm{Br}(S')$ is an isomorphism.*

Now suppose S is a smooth variety or orbifold over a field k , with function field $K = k(S)$. Then there is a description of the image of $\mathrm{Br}(S) \rightarrow \mathrm{Br}(K)$ in terms of residues. In case S is an orbifold, the role of the residue field at a point is played by the *residual gerbe*

$$\mathcal{G}_\xi \rightarrow \mathrm{Spec}(k(\xi))$$

at a point ξ of S (cf. [51, §11], [67, App. B]).

Proposition 2.2. *Let k be a field, S a smooth variety or orbifold over k with function field K , and n a positive integer that is invertible in k . Then there is a residue map from the n -torsion of the Brauer group of K to the direct sum of H^1 groups of residual gerbes \mathcal{G}_ξ at codimension 1 points $\xi \in S^{(1)}$, which fits into an exact sequence*

$$0 \rightarrow \mathrm{Br}(S)[n] \rightarrow \mathrm{Br}(K)[n] \rightarrow \bigoplus_{\xi \in S^{(1)}} H^1(\mathcal{G}_\xi, \mathbb{Z}/n\mathbb{Z}).$$

Proof. We already know that $\mathrm{Br}(S) \rightarrow \mathrm{Br}(K)$ is injective. Cohomological purity [11, §XVI.3], combined with the local-to-global spectral sequence

for cohomology with supports [59, §VI.5], yields exact sequences

$$\begin{aligned} \dots &\rightarrow H^{j-1}(S \setminus T, \mathbb{Z}/n\mathbb{Z}(i)) \rightarrow H^{j-2c}(T, \mathbb{Z}/n\mathbb{Z}(i-c)) \\ &\rightarrow H^j(S, \mathbb{Z}/n\mathbb{Z}(i)) \rightarrow H^j(S \setminus T, \mathbb{Z}/n\mathbb{Z}(i)) \rightarrow \dots \end{aligned}$$

for all i when $T \subset S$ is a closed substack of pure codimension c that is also smooth over k . We may assume, as in [17, Rmk. 4.7], that k is a perfect field, and as in [33] define the residue map and prove exactness. \square

Remark 2.3. When S is a smooth variety the exact sequence is well known, cf. [21], and may be extended with a further residue map as a complex (Bloch-Ogus complex)

$$\bigoplus_{\xi \in S^{(1)}} H^1(k(\xi), \mathbb{Z}/n\mathbb{Z}) \rightarrow \bigoplus_{\xi \in S^{(2)}} H^0(k(\xi), \mu_n^{-1}).$$

The complex with this additional residue map is exact when k is algebraically closed and S is a smooth projective rational surface.

2.4. Root stacks. We recall and describe basic properties of root stacks. Applied to a smooth variety and smooth divisor, the root stack construction produces an orbifold. The main function of a root stack is to remove ramification (e.g., of a Brauer class). In geometry, we find several advantages:

- We are able to relate smooth families over a root stack to flat families over the underlying scheme.
- The classification of smooth families may be more rigid, e.g., for conic bundles over a surface, the general ones correspond to Brauer classes at the generic point, while the smooth ones correspond to unramified Brauer classes.
- Other applications such as those requiring deformation theory also benefit from working with a proper Deligne-Mumford stack, instead of a quasi-projective variety.

Let S be a regular integral scheme, $D \subset S$ an effective divisor, and n a positive integer, invertible in the local rings of S . Then there is the corresponding *root stack* $\sqrt[n]{(S, D)}$, an integral locally Noetherian Deligne-Mumford stack, regular if and only if D is regular [20, §2], [1, App. B]. Above D is the *gerbe of the root stack* \mathcal{G}_D [20, Def. 2.4.4], an effective divisor on $\sqrt[n]{(S, D)}$ fitting into a commutative diagram

$$\begin{array}{ccc} \mathcal{G}_D & \longrightarrow & \sqrt[n]{(S, D)} \\ \downarrow & & \downarrow \\ D & \longrightarrow & S \end{array} \tag{2.1}$$

where the right-hand morphism is flat and is ramified over D (so the diagram is not cartesian unless D is empty) and restricts to an isomorphism over $S \setminus D$. The left-hand morphism makes \mathcal{G}_D into a μ_n -gerbe over D .

Remark 2.4. In the situation of Proposition 2.2, let ξ be the generic point of an irreducible regular divisor D . The ramification of the right-hand map in (2.1) leads us to expect a compatibility of residue maps with factor n playing the role of the ramification index, as explained by Saltman [69, Thm. 10.4] in the classical setting of discrete valuation rings and residue fields; so, for n -torsion classes, compatibility takes the form of a commutative diagram with residue maps as horizontal maps

$$\begin{array}{ccc} \mathrm{Br}(K)[n] & \longrightarrow & H^1(\mathcal{G}_\xi, \mathbb{Z}/n\mathbb{Z}) \\ \parallel & & \uparrow_0 \\ \mathrm{Br}(K)[n] & \longrightarrow & H^1(\mathrm{Spec}(k(\xi)), \mathbb{Z}/n\mathbb{Z}) \end{array}$$

and the zero map on the right. For the compatibility, we may replace S by $\mathrm{Spec}(\mathcal{O}_{S,\xi}^h)$, where $\mathcal{O}_{S,\xi}^h$ denotes the Henselization of the local ring of S at ξ ; we recall that passage to the residue field induces, then, an isomorphism on $H^1(-, \mathbb{Z}/n\mathbb{Z})$ [59, III.3.11(a)]. The compatibility is established by Lieblich [54, §3.2]: it suffices to treat just one n -torsion Brauer class having, as residue, any given class in the bottom right group, and such a Brauer class, obtained by pairing the class of a degree n cyclic étale algebra over $\mathcal{O}_{S,\xi}^h$ with a uniformizer, is shown to extend to the root stack.

Now let $D' \subset S$ be another effective divisor. The fiber product $\sqrt[n]{(S, D)} \times_S \sqrt[n]{(S, D')}$ of the two associated root stacks is the *iterated root stack* $\sqrt[n]{(S, \{D, D'\})}$ [20, Def. 2.2.4], with gerbes over respective divisors

$$\mathfrak{D} = \mathcal{G}_D \times_S \sqrt[n]{(S, D')} \quad \text{and} \quad \mathfrak{D}' = \sqrt[n]{(S, D)} \times_S \mathcal{G}_{D'}.$$

If D and D' are regular and intersect transversally, then $\sqrt[n]{(S, \{D, D'\})}$ is regular, and this remains valid with any number of divisors; we will speak of *rooted divisors* in an iterated root stack construction.¹ Let us suppose that $D \cap D'$ is nonempty; notice, the iterated root stack $\sqrt[n]{(S, \{D, D'\})}$ is different from the root stack $\sqrt[n]{(S, D \cup D')}$, which is singular. If we blow up $D \cap D'$ to obtain $\tilde{S} \rightarrow S$, then we have iterated root stack

¹A generalization, not used in this paper, is the case of a divisor D that étale locally may be written as a union of regular divisors meeting transversally, for which a construction by Matsuki and Olsson [56] based on log structures gives rise to a regular Deligne-Mumford stack, étale locally isomorphic to an iterated root stack.

$\sqrt[n]{(S, \{\tilde{D}, \tilde{D}', E\})}$ of the proper transforms and the exceptional divisor E , which is regular, while $\tilde{S} \times_S \sqrt[n]{(S, \{D, D'\})}$ is singular.

Zariski locally on S the divisor D is defined by the vanishing of a regular function f , and then $\sqrt[n]{(S, D)}$ is the quotient stack of

$$\mathrm{Spec}(\mathcal{O}_S[t]/(t^n - f))$$

by the scalar action of μ_n on t . The gerbe of the root stack is, Zariski locally, defined by the vanishing of t , and its points are just the points of D , with μ_n -stabilizer, i.e., above any point $x \in D$ with residue field $\kappa(x)$ the gerbe of the root stack takes the form of a classifying stack BG for $G = \mu_{n, \kappa(x)}$.

Often we consider flat families of projective varieties, or schemes, and we benefit by being able to study such families over a root stack, i.e., flat projective morphisms $P \rightarrow \sqrt[n]{(S, D)}$ of Deligne–Mumford stacks. The fiber over a point $s \in D$, where the gerbe of the root stack takes the form BG for $G = \mu_{n, \kappa(s)}$ as mentioned above, has the form $[P_s/G]$ for an action of G on a projective scheme P_s over $\kappa(s)$. Typically G acts with stabilizers (μ_n or subgroups of μ_n), and we will speak of the locus with μ_n -stabilizer for the appropriate closed subscheme of P_s .

2.5. Coarse moduli spaces. Let X be a Deligne–Mumford stack, and assume that X has *finite stabilizer*, meaning that projection to X from $X \times_{X \times X} X$ is a finite morphism. Then there is a *coarse moduli space*, an algebraic space Q with separated morphism $X \rightarrow Q$, universal for morphisms to algebraic spaces. For instance, X could be $[\mathrm{Spec}(A)/G]$ for some finite group G acting on affine scheme $\mathrm{Spec}(A)$, and then X has coarse moduli space $\mathrm{Spec}(A^G)$, where A^G denotes the G -invariant subring of A . The article by Keel and Mori [42] gives the classic treatment of coarse moduli space in the setting where X is locally of finite type over a locally Noetherian base scheme; this suffices for our purposes, and in this case Q has the same property and $X \rightarrow Q$ is proper and quasi-finite. The treatment without finiteness hypothesis is due to Rydh [68]. (These treatments allow, more generally, X to be an Artin stack with finite stabilizer.)

The construction of coarse moduli space shows that $X \rightarrow Q$ has the étale local (over Q) form $[\mathrm{Spec}(A)/G] \rightarrow \mathrm{Spec}(A^G)$, as described above. In fact, G may be taken to be the geometric stabilizer group at a point of X , and the assertion is valid on an étale neighborhood of the corresponding point of Q ; cf. [6, Lem. 3.4].

For the next statement we assume that X is *tame*, meaning that besides the finite stabilizer hypothesis we also require the order of the geometric

stabilizer group at any point $\text{Spec}(\Omega) \rightarrow X$ not to be divisible by the characteristic of Ω [65]. (This notion is also available for Artin stacks with finite stabilizer [2].) We suppose furthermore that X , and hence as well Q , is Noetherian. Then by [4, Thm. 10.3], pullback by $X \rightarrow Q$ identifies the category of vector bundles on Q with the full subcategory of vector bundles on X with trivial actions of geometric stabilizer groups at closed points.

The most relevant example of coarse moduli space for us is a root stack $\sqrt[n]{(S, D)} \rightarrow S$ (or an iterated root stack), and this is tame. The next result concerns schemes, *Fano over S* for a Noetherian scheme S , by which we mean the full subcategory of schemes over S , in which an object is a flat projective morphism $Z \rightarrow S$ with Gorenstein fibers such that $\omega_{Z/S}^\vee$ is relatively ample. The last condition is equivalent by [31, 4.7.1] to the ampleness of $\omega_{Z_s}^\vee$ on the fiber Z_s for every closed point $s \in S$. We may also take S to be an algebraic space, in which case Z is also allowed to be an algebraic space. We may allow S to be a DM stack, in which case Z is also a DM stack. But since stacks form a 2-category, we need to take care to obtain an ordinary category: a morphism from $f: Z \rightarrow S$ to $f': Z' \rightarrow S$ is defined to be an equivalence class of pairs (φ, α) where $\varphi: Z \rightarrow Z'$ is a morphism and $\alpha: f \Rightarrow f' \circ \varphi$ is a 2-morphism, with $(\varphi, \alpha) \sim (\tilde{\varphi}, \tilde{\alpha})$ when there exists a (necessarily unique) 2-morphism $\beta: \varphi \Rightarrow \tilde{\varphi}$, such that $f'(\beta(z)) \circ \alpha(z) = \tilde{\alpha}(z)$ for every object z of Z .

Proposition 2.5. *Let X be a tame Noetherian DM stack with coarse moduli space $X \rightarrow Q$. Then the category of schemes, Fano over Q , is equivalent, by base change, to the full subcategory of DM stacks, Fano over X , with trivial actions of geometric stabilizer groups on fibers at closed points.*

Proof. The universal property of the coarse moduli space tells us that base change by $X \rightarrow Q$ is a fully faithful functor from the category of schemes, Fano over Q , to the category of DM stacks, Fano over X .

To an object $f: Z \rightarrow X$ of the category of DM stacks, Fano over X , there is an associated quasi-coherent sheaf of graded \mathcal{O}_X -algebras $\bigoplus_{n \geq 0} f_*((\omega_{Z/X}^\vee)^n)$. The graded components are coherent and for sufficiently large n are locally free with formation commuting with base change to (geometric) fibers. If the actions of geometric stabilizer groups at closed points is trivial, then $f_*((\omega_{Z/X}^\vee)^n)$ for $n \gg 0$ satisfies the condition stated above to descend to a locally free coherent sheaf on Q . The algebra structure descends as well, and by applying Proj we obtain an algebraic space Y , Fano over Q , which upon base change to X recovers $Z \rightarrow X$ up to an isomorphism. \square

Remark 2.6. As a consequence, the algebraic space Y , Fano over Q that we obtain in the proof of Proposition 2.5, is the coarse moduli space of Z .

2.6. Étale local uniqueness in smooth families. In certain geometric situations such as fibrations in projective spaces, a smooth family has a unique étale local isomorphism type. The same holds G -equivariantly, when G is a finite group whose order is not divisible by the residue characteristic.

Lemma 2.7. *Let A be a Henselian ring with residue field κ , and X a scheme, smooth over $\text{Spec}(A)$. Let G be a finite group, whose order is invertible in κ , with compatible actions on X and on A and trivial action on κ , and x a κ -point of the fiber X_κ over the closed point of $\text{Spec}(A)$ that is fixed for the group action. Then X admits a G -invariant A -point that specializes to x .*

Proof. Since A and hence as well A^G is Henselian, any finite-dimensional κ -vector space with G -action extends to a free finite-rank A^G -module with G -action, by Henselian lifting [32, 18.5.17], applied to the scheme of homomorphisms $G \rightarrow GL_n$ (which is smooth since the order of G is invertible in κ), where n is the dimension of the given κ -vector space.

Replacing, as needed, X by intersections of translates under G of an affine neighborhood of x , we may suppose that $X = \text{Spec}(B)$ is affine. Let \mathfrak{m} denote the maximal ideal corresponding to x . The A -linear map

$$\mathfrak{m} \rightarrow \Omega_{X_\kappa, x}^1$$

(sending f to the Kähler differential df , over κ) is surjective and G -equivariant. There is therefore a lift of an appropriate κ -basis of $\Omega_{X_\kappa, x}^1$ to elements of \mathfrak{m} whose A^G -linear span is invariant under the G -action. According to [32, 17.16.3(i)] (really, its proof), the lifts define a G -invariant subscheme $W \subset X$ that is étale over $\text{Spec}(A)$ at x . Since A is Henselian, there is a unique G -invariant A -point of W that specializes to x . \square

Lemma 2.8. *Let A be a Henselian ring with residue field κ , and X and Y schemes, smooth and projective over $\text{Spec}(A)$. Let G be a finite group, whose order is invertible in κ , with compatible actions on X , on Y , and on A , with trivial action on κ . Let $\varphi: X_\kappa \rightarrow Y_\kappa$ be a G -equivariant isomorphism of fibers. If the tangent bundle T_{X_κ} satisfies*

$$H^1(X_\kappa, T_{X_\kappa}) = 0, \tag{2.2}$$

then there exists a G -equivariant isomorphism $X \rightarrow Y$ over $\text{Spec}(A)$ that specializes to φ .

Proof. There is a scheme $\text{Isom}_A(X, Y)$, parametrizing isomorphisms of fibers [34, §4c], with G -action, which by (2.2) is smooth over $\text{Spec}(A)$ [34, Cor. 5.4]. We conclude by Lemma 2.7. \square

2.7. Components in families. The irreducible components of fibers in a flat family with geometrically reduced fibers may be accessed by excluding the relative singular locus and forming the sheaf of connected components. If $\pi: X \rightarrow S$ is a finitely presented smooth morphism of schemes, then geometric components of fibers determine an étale equivalence relation on X over S whose algebraic space quotient $\pi_0(X/S)$ is étale over S and is characterized as factorizing π ,

$$X \rightarrow \pi_0(X/S) \rightarrow S,$$

such that $X \rightarrow \pi_0(X/S)$ is surjective with geometrically connected fibers [51, 6.8]. There are easy examples for which $\pi_0(X/S)$ is not a scheme, e.g., the sum of squares of coordinates $\mathbb{A}_{\mathbb{R}}^2 \setminus \{(0, 0)\} \rightarrow \mathbb{A}_{\mathbb{R}}^1$. Formation of $\pi_0(X/S)$ commutes with base change by an arbitrary morphism $S' \rightarrow S$.

A subset of X , which is the pre-image of a subset of $\pi_0(X/S)$, will be called *component-saturated*.

Lemma 2.9. *Let $\pi: X \rightarrow S$ be a proper flat finitely presented pure-dimensional morphism of schemes, such that for every $s \in S$ the smooth locus of the fiber X_s is dense in X_s . Then the morphism $\pi_0(X^{\text{sm}}/S) \rightarrow S$ is universally closed, where X^{sm} denotes the relative smooth locus of π . If, moreover, the geometric fibers of π all have d irreducible components for some integer d , then $\pi_0(X^{\text{sm}}/S) \rightarrow S$ is finite étale of degree d .*

Proof. The second assertion follows from the first by the general fact that a finitely presented universally closed étale morphism $T \rightarrow S$ with fibers all of the same degree is finite. This comes down to topological properties of the relative diagonal (it is an open immersion since $T \rightarrow S$ is étale, and we need to show that it is closed as well), so the verification is reduced by [32, 1.10.1] to the case that S is Spec of a local ring, and by [32, 2.6.2], to the case of a strictly Henselian ring. Each of the d pre-images of the closed point then lifts to a section. The sections cover T , since the image of their complement is closed but does not contain the closed point of S . So, T is the disjoint union of the sections.

Now suppose that $S = \text{Spec}(A)$, where A is a local ring. Let $z \in X$ be a maximal point of a fiber of π and Z its closure in X . We claim that $\pi(Z \cap X^{\text{sm}})$ contains the closed point of S . Indeed, by semicontinuity of fiber dimension [32, 13.1.5], Z contains a maximal point of the fiber over the closed point, which by hypothesis lies in X^{sm} .

Let $n \geq 0$. In a proper flat family of schemes of relative dimension n over a Noetherian base scheme, the condition on fibers to have singular

locus of dimension less than n is open by semicontinuity of fiber dimension. With the local form of semicontinuity of fiber dimension [32, 13.1.3] (semicontinuity of $w \mapsto \min \dim(\pi^{-1}(\pi(w)) \cap U)$ where the minimum is over open U containing w) and flatness (particularly, [32, 2.3.4]), we see that the condition to have pure dimension n is closed. We reduce the verification of the first assertion immediately to the case when S is affine and reduced, and with the preceding observations and [32, 8.10.5, 11.2.7] to the case when S is Noetherian. Then, as noted in the proof of [46, Prop. 2.2], the claim implies the assertion. (In fact, the case A is a DVR suffices.) \square

Remark 2.10. With the notation of Lemma 2.9, any component-saturated closed subset $W \subset X^{\text{sm}}$ has the property that formation of the closure \overline{W} in X commutes with base change by an arbitrary morphism $S' \rightarrow S$. Since an equality of sets may be verified fiberwise, the assertion reduces to the case $S' = \text{Spec}(\kappa(s))$, where $\kappa(s)$ is the residue field of a point $s \in S$. We may suppose π has constant fiber dimension n . For $x \in \overline{W} \cap \pi^{-1}(s)$, semicontinuity of fiber dimension yields $\dim(\pi^{-1}(s) \cap U) = n$ for all open $U \subset \overline{W}$ containing x . Hence there is a maximal point of $\pi^{-1}(s)$ that lies in W (equivalently, lies in \overline{W}) and specializes to x .

3. CONIC BUNDLES

In this section we present an ingredient from the Sarkisov program, which gives a description of conic bundles with regular discriminant divisor, using the language of root stacks.

By a conic bundle over a regular scheme S , in which 2 is invertible in the local rings, we mean a flat projective morphism $\pi: X \rightarrow S$ with smooth generic fiber over every irreducible component of S , such that the fiber over every point of S is isomorphic to a possibly singular, possibly non-reduced conic in \mathbb{P}^2 , i.e., the zero-locus of a nontrivial ternary quadratic form. We say that a conic bundle $\pi: X \rightarrow S$ has *mild degeneration* if all fibers are reduced.

Proposition 3.1. *Let S be a regular scheme, such that 2 is invertible in the local rings of S , and let $D \subset S$ be a regular divisor. Then the operations described below identify, up to unique isomorphism:*

- *smooth \mathbb{P}^1 -fibrations over $\sqrt{(S, D)}$ having nontrivial μ_2 -actions on all fibers over the gerbe of the root stack, with*
- *conic bundles $\pi: X \rightarrow S$ such that X is regular and the fiber $\pi^{-1}(s)$ over a point $s \in S$ is singular if and only if $s \in D$; all such conic bundles have mild degeneration.*

Given a smooth \mathbb{P}^1 -fibration $P \rightarrow \sqrt{(S, D)}$ with nontrivial μ_2 -actions on all fibers over the gerbe of the root stack, a conic bundle $\pi: X \rightarrow S$ with mild degeneration is obtained by

- blowing up the locus with μ_2 -stabilizer,
- contracting the “middle components” of fibers over D , and
- descending to S .

Given a conic bundle $\pi: X \rightarrow S$ we obtain the associated smooth \mathbb{P}^1 -fibration over $\sqrt{(S, D)}$ (up to unique isomorphism) by

- pulling back to $\sqrt{(S, D)}$,
- blowing up the relative singular locus over the gerbe of the root stack, and
- contracting the “end components” of fibers over D .

Proposition 3.1 deals with flat projective families of reduced curves with at most nodes as singularities. This is the setting of (pre)stable curves, for which there are known constructions to perform the contractions indicated in the statement of Proposition 3.1. Contracting the “middle components” of a flat family of at-most-3-component genus 0 prestable curves $\pi: C \rightarrow T$ is achieved by the relative anticanonical model $\text{Proj}(\bigoplus_{m \geq 0} \pi_*((\omega_{C/T}^\vee)^m))$. The setting where “end components” are to be contracted is one of families with only 1- and 3-component genus 0 prestable curves, and étale locally there exists a section σ which avoids the end components of 3-component fibers; then $\text{Proj}(\bigoplus_{m \geq 0} \pi_*({\mathcal O}_C(\sigma)^m))$ gives the desired contraction, and although sections exist only étale locally, the construction may be performed globally over the base, as described in [46].

Proof. Let $P \rightarrow \sqrt{(S, D)}$ be a smooth \mathbb{P}^1 -fibration, such that μ_2 acts nontrivially on P_s for every $s \in D$ in the notation from the discussion of stabilizers from §2.4. Let $\text{Spec}(A)$ be an affine neighborhood in S of a point $s \in D$, on which D is defined by the vanishing of a function $f \in A$. Setting

$$A' := A[t]/(t^2 - f),$$

then, we have an open substack of $\sqrt{(S, D)}$ isomorphic to

$$[\text{Spec}(A')/\mu_2],$$

where μ_2 acts by scalar multiplication on t . We claim that there is an étale neighborhood $\text{Spec}(B) \rightarrow \text{Spec}(A)$ of s such that for the rank 2 vector bundle and projectivization

$$V := [\text{Spec}(A'[u, v])/\mu_2] \quad \text{and} \quad \widehat{P} := \mathbb{P}(V),$$

where μ_2 acts by scalar multiplication on u and trivially on v , we have

$$\mathrm{Spec}(B) \times_S P \cong \mathrm{Spec}(B) \times_S \widehat{P}.$$

Equivalently,

$$\mathrm{Spec}(A^{sh}) \times_S P \cong \mathrm{Spec}(A^{sh}) \times_S \widehat{P},$$

where A^{sh} is a strict Henselization of the local ring of $\mathrm{Spec}(A)$ at s . By [32, 18.8.10], we have

$$A^{sh} \otimes_A A' \cong A'^{sh},$$

where A'^{sh} is an analogous strict Henselization of A' . So, it suffices to verify that

$$\mathrm{Spec}(A'^{sh}) \times_{\sqrt{(S,D)}} P \quad \text{and} \quad \mathrm{Spec}(A'^{sh}) \times_{\sqrt{(S,D)}} \widehat{P}$$

are equivariantly isomorphic. By Lemma 2.8 it is enough to check that the respective fibers over the closed point of $\mathrm{Spec}(A'^{sh})$ are equivariantly isomorphic, and this is so by the hypothesis of nontrivial μ_2 -action on fibers of P over the gerbe of the root stack. In particular, the stabilizer locus of P is a degree 2 étale representable cover of the gerbe of the root stack.

In summary, the given \mathbb{P}^1 -fibration $P \rightarrow \sqrt{(S,D)}$ is, étale locally over S , isomorphic to the \mathbb{P}^1 -fibration $\widehat{P} \rightarrow \sqrt{(S,D)}$ given by a coordinate action of μ_2 on \mathbb{P}^1 .

The blow-up P_1 of the locus of P with μ_2 -stabilizer therefore has geometric fibers over points of D with three irreducible components and nontrivial μ_2 -action only on the “middle components”. After contracting “middle components” to P_2 , the geometric fibers have two irreducible components over points of D and trivial μ_2 -action. By Proposition 2.5, $P_2 \rightarrow \sqrt{(S,D)}$ descends to a conic bundle $P_3 \rightarrow S$. To see that P_1 is the blow-up of P_2 along the relative singular locus over the gerbe of the root stack, we use the fact that the constructions described in the statement are compatible with base change by an étale morphism to S and carry out the constructions with \widehat{P} :

$$\begin{aligned} \widehat{P}_1 &= [\mathrm{Proj}(A'[p, q, r, w]/(pr - tq^2, pw - t^2qr, qw - tr^2))/\mu_2], \\ \widehat{P}_2 &= [\mathrm{Proj}(A'[x, y, z]/(xz - t^2y^2))/\mu_2], \\ \widehat{P}_3 &= \mathrm{Proj}(A[x, y, z]/(xz - fy^2)). \end{aligned}$$

Since \widehat{P}_3 is regular, so is P_3 . The steps, performed in reverse starting from P_3 , yield P up to unique isomorphism.

A conic bundle $X \rightarrow S$ with X regular and singular fibers precisely over points of D has, locally, defining equation $fx^2 + ay^2 + bz^2 = 0$ for units a and b , as we see using diagonalizability of a quadratic form over

$\mathcal{O}_{S,s}$, first with s the generic point of a component of D and then with arbitrary $s \in D$. (Such an observation for quadric bundles is given in [13, Prop. 1.2.5].) So, étale locally, there is an isomorphism with \widehat{P}_3 , and the construction may be carried out to yield a smooth \mathbb{P}^1 -bundle over $\sqrt{(S,D)}$. From this, the construction of blowing up, contracting, and descending to S recovers $X \rightarrow S$ up to unique isomorphism. \square

Remark 3.2. The Sarkisov program establishes the existence of good models of conic bundles over arbitrary varieties [70], via classical birational geometry. A more general version of this result may be obtained through the use of root stacks. Details are presented in [64].

4. BRAUER-SEVERI SURFACE BUNDLES

In the following sections we prove Theorem 1.2 using root stacks.

Definition 4.1. Let S be a regular scheme, in which 3 is invertible in the local rings. A *Brauer-Severi surface bundle* over S is a flat projective morphism $\pi: X \rightarrow S$ with smooth generic fiber over every irreducible component of S , such that the fiber over every geometric point of S is isomorphic to one of the following:

- \mathbb{P}^2 ,
- the union of three Hirzebruch surfaces \mathbb{F}_1 , meeting transversally, such that any pair of them meets along a fiber of one and the (-1) -curve of the other,
- an irreducible scheme whose underlying reduced subscheme is isomorphic to the cone over a twisted cubic curve.

If only fibers of the first two types appear, then we say that the Brauer-Severi surface bundle has *mild degeneration*.

Remark 4.2. For the fibers X_s appearing in a Brauer-Severi surface bundle with mild degeneration, the dual $\omega_{X_s}^\vee$ of the dualizing sheaf is very ample and we have $h^0(X_s, \omega_{X_s}^\vee) = 10$. This is standard for \mathbb{P}^2 and may be verified with a short computation for unions of three Hirzebruch surfaces, using that the restriction of $\omega_{X_s}^\vee$ to each component is the class of the (-1) -curve plus two fibers.

Definition 4.3. Let k be a field of characteristic different from 3. An action of μ_3 on \mathbb{P}_k^2 is said to be *balanced* if, after passing to an algebraic closure \bar{k} of k and making a change of projective coordinates, the action is $\mu_3 \times \mathbb{P}_{\bar{k}}^2 \rightarrow \mathbb{P}_{\bar{k}}^2$,

$$(\alpha, x : y : z) \mapsto (\alpha x : \alpha^2 y : z).$$

We note that an action of μ_3 on \mathbb{P}_k^2 is balanced if and only if the fixed-point locus has dimension zero.

We formulate an analogue of Proposition 3.1. In the statement, degree of a component refers to the degree of the anticanonical class of the component.

Proposition 4.4. *Let S be a regular scheme, such that 3 is invertible in the local rings of S , and let $D \subset S$ be a regular divisor. Then the operations described below identify, up to unique isomorphism:*

- smooth \mathbb{P}^2 -fibrations over $\sqrt[3]{(S, D)}$ having balanced μ_3 -actions on fibers over the gerbe of the root stack, with
- Brauer-Severi surface bundles $\pi: X \rightarrow S$ such that X is regular and the fiber $\pi^{-1}(s)$ over a point $s \in S$ is singular if and only if $s \in D$; all such bundles have mild degeneration.

Given a smooth \mathbb{P}^2 -fibration $P \rightarrow \sqrt[3]{(S, D)}$ with balanced μ_3 -actions on fibers over the gerbe of the root stack, P has distinguished loci

- P_μ : locus with nontrivial μ_3 -stabilizer,
- P_λ : union of lines on the geometric \mathbb{P}^2 -fibers joining points of P_μ .

On the blow-up $B\ell_{P_\mu} P$ of P we have a distinguished locus

- P_c : the union of codimension 2 components of the stabilizer locus.

On the second blowup $B\ell_{P_c}(B\ell_{P_\mu} P)$ we have the locus

- \tilde{P}_λ : the proper transform of P_λ .

Then a Brauer-Severi surface bundle $\pi: X \rightarrow S$ with mild degeneration results by

- contracting first the degree 8 components of fibers over D of

$$B\ell_{\tilde{P}_\lambda}(B\ell_{P_c}(B\ell_{P_\mu} P)),$$

- contracting the degree 3 components,
- descending to S .

Given a Brauer-Severi surface bundle $\pi: X \rightarrow S$ we obtain the associated smooth \mathbb{P}^2 -fibration over the root stack $\sqrt[3]{(S, D)}$ (up to unique isomorphism) by

- pulling back to $\sqrt[3]{(S, D)}$,
- blowing up the relative singular locus of the relative singular locus over the gerbe of the root stack,
- blowing up the relative singular locus over the gerbe of the root stack, and
- contracting the high-degree (≥ 7) components above the gerbe of the root stack.

The constructions in Proposition 4.4 are more complicated than the corresponding constructions for conic bundles. Their justification relies on machinery presented in an Appendix on birational contractions.

The proof, given in the following sections, has several ingredients:

- the forwards construction, which produces a mildly degenerating Brauer-Severi surface bundle out of a smooth of \mathbb{P}^2 -fibration over the root stack, is carried out using explicit birational contractions (Section 5);
- the reverse construction, starting with a mildly degenerating Brauer-Severi surface bundle, is carried out with similar techniques (Section 6);
- the verification that every Brauer-Severi surface bundle $\pi: X \rightarrow S$, such that X is regular and precisely the fibers over a regular divisor on S are singular, has mild degeneration, based on a combination of the forwards construction and purity for the Brauer group (Section 7);

5. FORWARDS CONSTRUCTION

In this section we carry out the forwards construction of Proposition 4.4.

Let $P \rightarrow \sqrt[3]{(S, D)}$ be a smooth \mathbb{P}^2 -fibration, such that the μ_3 -action on P_s for every $s \in D$ is balanced. As in the proof of Proposition 3.1, there is an affine étale neighborhood $\text{Spec}(B) \rightarrow S$ of a point $s \in D$, where the pre-image of D is principal, say, defined by the vanishing of $f \in B$, and setting

$$B' := B[t]/(t^3 - f), \quad V := [\text{Spec}(B'[u, v, w])/\mu_3], \quad \widehat{P} := \mathbb{P}(V),$$

we have an isomorphism

$$\text{Spec}(B) \times_S P \cong \widehat{P}. \tag{5.1}$$

Here, μ_3 acts on t by scalar multiplication, on u by scalar multiplication squared, on v , and trivially on w . In particular, P_μ is a degree 3 étale representable cover of the gerbe of the root stack, and after blowing up P_c is smooth over the gerbe of the root stack, consisting of a line in every \mathbb{P}^2 fiber of the exceptional divisor, and \tilde{P}_λ is also smooth over the gerbe of the root stack, consisting geometrically of three pairwise disjoint (-2) -curves in every component of fibers over the gerbe of the root stack, isomorphic to the minimal resolution of the singular cubic surface of type $3A_2$ [19].

At the successive blow-ups we have divisors at our disposal as described below; for the description of combinations that are relatively ample over S we use the fact about ampleness mentioned in §2.5 in combination with the equivalence [31, 2.6.2] of ampleness of a line bundle on a projective scheme with ampleness of its restriction to every irreducible component, to characterize relatively ample line bundles. We adopt the convention

that the same notation will be used for a divisor or line bundle and its pullbacks under the morphisms described below. But in this section, tilde ($\tilde{\cdot}$), as in \tilde{E}_1 , always indicates proper transform. We let ω^\vee denote the relative anticanonical bundle of P over $\sqrt[3]{(S, D)}$.

- We analyze the first blow-up

$$B\ell_{P_\mu} P \rightarrow P.$$

In addition to ω^\vee , we have the exceptional divisor E_1 ; on geometric fibers over the gerbe of the root stack, E_1 consists of three components E_u, E_v, E_w , each isomorphic to \mathbb{P}^2 , attached along three pairwise disjoint exceptional divisors d_u, d_v, d_w of a component Σ_1 isomorphic to a degree 6 del Pezzo surface (DP6), with the remaining exceptional divisors $\lambda_{uv}, \lambda_{vw}, \lambda_{uw} \subset \Sigma_1$ contained in the proper transform of P_λ . The intersection numbers are:

ω^\vee	d_u	λ_{vw}
	0	3
E_1	-1	2

The projection formula supplies the first row. Entries in the second row are intersection numbers on DP6 with $d_u + d_v + d_w$.

Multiples of $\alpha\omega^\vee - E_1$ for $\alpha \in \mathbb{Q}$, $\alpha > 2/3$ are relatively ample.

- We analyze the second blowup

$$B\ell_{P_c}(B\ell_{P_\mu} P) \rightarrow B\ell_{P_\mu} P.$$

Let E_2 be its exceptional divisor. With notation as above and components $\ell_u \subset E_u$, etc., of the stabilizer locus P_c , we have the exact sequence of normal bundles

$$0 \rightarrow N_{P_c/E_1}|_{\ell_u} \rightarrow N_{P_c/B\ell_{P_\mu} P}|_{\ell_u} \rightarrow N_{E_1/B\ell_{P_\mu} P}|_{\ell_u} \rightarrow 0$$

with

$$\deg(N_{\ell_u/E_u}) = 1 \quad \text{and} \quad \deg(N_{E_u/B\ell_{P_\mu} P}|_{\ell_u}) = -1$$

(and similarly for ℓ_v and ℓ_w). On geometric fibers over the gerbe of the root stack, E_2 consists of three components F_u, F_v, F_w , each isomorphic to a Hirzebruch surface \mathbb{F}_2 , glued along (-2) -curves c_u , respectively, c_v, c_w , to \tilde{E}_u , respectively, \tilde{E}_v, \tilde{E}_w , and along a fiber $f_{2,u}$ (respectively, $f_{2,v}, f_{2,w}$) to Σ_2 , the blow-up of Σ_1 at three points; note that Σ_2 is isomorphic to the minimal resolution of the singular cubic surface of type $3A_2$. The class of the exceptional divisor E_2 , restricted to F_u , etc., is $\mathcal{O}_{\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1))}(-1)$. The further restriction to c_u , etc., is $\mathcal{O}_{\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(1) \oplus 0)}(-1) \cong \mathcal{O}_{\mathbb{P}^1}(1)$ and

to $f_{2,u}$, etc., is isomorphic to $\mathcal{O}_{\mathbb{P}^1}(-1)$; these explain the last two entries in the third row:

	\tilde{d}_u	$\tilde{\lambda}_{vw}$	c_u	$f_{2,u}$
ω^\vee	0	3	0	0
E_1	-1	2	-1	0
E_2	1	1	1	-1

Intersection numbers on Σ_2 give the first two entries in the third row.

Multiples of $\beta(\alpha\omega^\vee - E_1) - E_2$ with

$$\alpha > (1/3)(2 + 1/\beta) \quad \text{and} \quad \beta > 1$$

are relatively ample.

- We analyze the third blowup

$$B\ell_{\tilde{P}_\lambda}(B\ell_{P_c}(B\ell_{P_\mu} P)) \rightarrow B\ell_{P_c}(B\ell_{P_\mu} P).$$

Let E_3 be its exceptional divisor. On geometric fibers over the gerbe of the root stack E_3 consists of components Q_{uv} , Q_{vw} , Q_{uw} , fibered over $\tilde{\lambda}_{uv}$, etc. With an exact sequence of normal bundles, as in the second blowup, we see that each component is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$, with $\mathcal{O}(E_3)$ of degree -2 along ruling sections and -1 along fibers, and is glued along a fiber to the exceptional divisor e_u of a blow-up of \tilde{E}_u , etc., along another fiber to the exceptional divisor f_u of a blow-up of F_u , etc., and along a ruling section (in the table, $\tilde{\lambda}_{vw}$, etc.) to $\Sigma_3 \cong \Sigma_2$.

	\tilde{d}_u	e_u	$\tilde{\lambda}_{vw}$	c_u	$\tilde{f}_{2,u}$	f_u
ω^\vee	0	0	3	0	0	0
E_1	-1	0	2	-1	0	0
E_2	1	0	1	1	-1	0
E_3	1	-1	-2	0	1	-1

Multiples of $\gamma(\beta(\alpha\omega^\vee - E_1) - E_2) - E_3$ with

$$\alpha > \frac{2}{3} + \frac{1}{3\beta} - \frac{2}{3\beta\gamma}, \quad \beta > 1 + \frac{1}{\gamma}, \quad \text{and} \quad \gamma > 1$$

are relatively ample.

The degree 8 components of geometric fibers over the root stack \tilde{E}_u , \tilde{E}_v , \tilde{E}_w , Q_{uv} , Q_{vw} , Q_{uw} comprise the divisor

$$\tilde{E}_1 \cup E_3$$

in the class of $E_1 - E_2 + E_3$, with evident contraction ψ to a stack Z , isomorphic to each of the intersections $\tilde{E}_1 \cap E_2$, $\tilde{E}_1 \cap E_3$, $E_2 \cap E_3$. Proposition A.9 is applicable to the contraction $\psi: \tilde{E}_1 \cup E_3 \rightarrow Z$ with

the relatively ample line bundle associated with (an integral multiple of) $\gamma(\beta(\alpha\omega^\vee - E_1) - E_2) - E_3$ for

$$\alpha\beta > 1, \quad (1-\alpha)\beta < 1, \quad (2\alpha-1)\beta < 1, \quad \gamma = \frac{1}{(1-\alpha)\beta},$$

and value of m , coefficient appearing in the twist of the relatively ample line bundle by $m(E_1 - E_2 + E_3)$, equal to (the same integral multiple of) $(\alpha\beta-1)/(1-\alpha)\beta$. We obtain a contraction

$$\nu: B\ell_{\tilde{P}_\lambda}(B\ell_{P_e}(B\ell_{P_\mu} P)) \rightarrow P_1,$$

restricting to $\psi: \tilde{E}_1 \cup E_3 \rightarrow Z \subset P_1$, with $\mathcal{I}_Z/\mathcal{I}_Z^2$ locally free of rank 3.

By Remark A.11, the effective Cartier divisor $E_1 + E_3$, viewed as invertible sheaf with global section, pushes forward under ν to a Cartier divisor on P_1 that we denote by Ψ . Over $\text{Spec}(B)$, where we have an isomorphism (5.1) and therefore individually defined divisors \tilde{E}_u , etc., further Cartier divisors arise by pushing forward

$$\hat{G}_u := 2\tilde{E}_u + \tilde{E}_v + 3\tilde{F}_u + 2Q_{uv} + Q_{uw}$$

and the analogously defined \hat{G}_v and \hat{G}_w to

$$\hat{P}_1 := \text{Spec}(B) \times_S P_1.$$

So the union of the blown up \mathbb{F}_2 components of \hat{P}_1 over the gerbe of the root stack is a Cartier divisor, as is 3 times each individual blown up \mathbb{F}_2 component. Since ν is a birational contraction of a normal scheme, P_1 is normal.

We claim, P_1 has A_2 -singularities along Z . This is a local assertion, so we may work over \hat{P}_1 , which is a quotient stack by μ_3 . We use the language of schemes (with μ_3 -action) in the following argument. Fix a closed point $\hat{r} \in \hat{Z}$ over $s' \in \text{Spec}(B')$, corresponding to some $\mathfrak{m}' \subset B'$, over a given $s \in D$. Let $n = \dim(B'_{\mathfrak{m}'})$, and let t, g_1, \dots, g_{n-1} be a regular system of parameters, i.e., elements that form a basis of $\mathfrak{m}'/\mathfrak{m}'^2$. If $\hat{\mathfrak{n}}$ denotes the maximal ideal of the local ring $\mathcal{O}_{\hat{P}_1, \hat{r}}$, where the residue field of \hat{r} is a finite extension ℓ of the residue field k of s' , then since \hat{Z} is regular there is the exact sequence of vector spaces over ℓ [32, 16.9.13]

$$0 \rightarrow \mathcal{I}_{\hat{Z}}/\hat{\mathfrak{n}}\mathcal{I}_{\hat{Z}} \rightarrow \hat{\mathfrak{n}}/\hat{\mathfrak{n}}^2 \rightarrow \hat{\mathfrak{n}}/(\mathcal{I}_{\hat{Z}} + \hat{\mathfrak{n}}^2) \rightarrow 0, \quad (5.2)$$

from which we learn

$$\dim(\hat{\mathfrak{n}}/\hat{\mathfrak{n}}^2) = n + 3.$$

As well, by Proposition A.9, we have the compatibility of the contraction $\hat{\nu}$ with base change, e.g., via $\{s'\} \rightarrow \text{Spec}(B')$, and this way we learn that the fiber of \hat{P}_1 over s' consists of a singular cubic surface of type $3A_2$ joined

to three components, each isomorphic to the blow-up of \mathbb{F}_2 at one point. In particular,

$$\dim(\hat{\mathfrak{n}}/(\mathfrak{m}'\mathcal{O}_{\widehat{P}_1,\hat{r}} + \hat{\mathfrak{n}}^2)) = \begin{cases} 4, & \text{if 3 components meet at } \hat{r}, \\ 3, & \text{otherwise.} \end{cases}$$

In both cases, we let x and y be local defining equations of the Cartier divisors on \widehat{P}_1 , each consisting (on fibers) of one blown up \mathbb{F}_2 component with multiplicity 3, and z , of their union with multiplicity 1; in the case 3 components meet at \hat{r} we additionally let q be a local defining equation for the singular cubic surface divisor component.

We have, after adjusting by units,

$$xy = z^3$$

and, in the two cases,

$$t = qz \in \hat{\mathfrak{n}}^2, \quad \text{respectively,} \quad t = z.$$

The kernel

$$(\mathfrak{m}'\mathcal{O}_{\widehat{P}_1,\hat{r}} + \hat{\mathfrak{n}}^2)/\hat{\mathfrak{n}}^2$$

of the canonical homomorphism $\hat{\mathfrak{n}}/\hat{\mathfrak{n}}^2 \rightarrow \hat{\mathfrak{n}}/(\mathfrak{m}'\mathcal{O}_{\widehat{P}_1,\hat{r}} + \hat{\mathfrak{n}}^2)$ is generated by \mathfrak{m}' . So the kernel is, in the two cases, of dimension $n - 1$ with basis g_1, \dots, g_{n-1} , respectively, of dimension n with basis t, g_1, \dots, g_{n-1} .

Suppose, first, \hat{r} is the point where three components come together. We know, modulo q , the point \hat{r} is an A_2 -singular point of the cubic surface component, with $\hat{\mathfrak{n}}/(\mathfrak{m}'\mathcal{O}_{\widehat{P}_1,\hat{r}} + q\mathcal{O}_{\widehat{P}_1,\hat{r}} + \hat{\mathfrak{n}}^2)$ of dimension 3, spanned by x, y , and z . So q, x, y , and z span $\hat{\mathfrak{n}}/(\mathfrak{m}'\mathcal{O}_{\widehat{P}_1,\hat{r}} + \hat{\mathfrak{n}}^2)$; they define a morphism to $\mathbb{A}_{B'}^4$ which is unramified at \hat{r} and hence by [32, 18.4.7] restricts and lifts to a closed immersion of a neighborhood $\text{Spec}(C)$ of \hat{r} in an étale affine neighborhood $\text{Spec}(\Delta)$ of the point 0 over s' in $\mathbb{A}_{B'}^4$; let $\delta_q, \delta_x, \delta_y, \delta_z \in \Delta$ denote the coordinate functions from $\mathbb{A}_{B'}^4$. Then, setting $\tilde{\Delta} := \Delta/(t - \delta_q\delta_z)$, we see that $\Delta \rightarrow C$ factors through

$$\tilde{\Delta}/(\delta_x\delta_y - \delta_z^3). \tag{5.3}$$

The element $\delta_x\delta_y - \delta_z^3$ is irreducible in the local ring of $\tilde{\Delta}$ at the point 0 over s' , so the localization of (5.3) is an integral domain, and we have found the desired local form of the singularity at \hat{r} .

Now suppose \hat{r} is a point of the component \mathbb{P}^1 where two blown up \mathbb{F}_2 components meet. The Cartier divisor defined by x , restricted to the other blown up \mathbb{F}_2 component, defines \mathbb{P}^1 with multiplicity 1, and the same is true with y and the first blown up \mathbb{F}_2 component. It follows that x, y , and a polynomial of degree $[\ell : k]$ in a local parameter p for \mathbb{P}^1 span

$\hat{\mathfrak{n}}/(\mathfrak{m}'\mathcal{O}_{\hat{P}_1, \hat{r}} + \hat{\mathfrak{n}}^2)$. We argue as above, with morphism to $\mathbb{A}_{B'}^3$ determined by x, y , and p and $\Delta \rightarrow C$ factoring through $\Delta/(\delta_x\delta_y - t^3)$.

On P_1 we have Cartier divisors $\Psi \cup \nu(\Sigma_3)$ and $\bar{\Psi}$. Hence $\nu(\Sigma_3)$ is a Cartier divisor, which becomes linearly equivalent to $-\Psi$ after base change to any affine étale chart of $\sqrt[3]{(S, D)}$ where the pre-image of the gerbe of the root stack is principal. We apply Proposition A.9 with relatively ample $\mathcal{O}_{P_1}(1)$ and m taken so that the twist by $m\nu(\Sigma_3)$ has degree 0 on $\tilde{f}_{2,u}$. Thus we obtain a contraction

$$\rho: P_1 \rightarrow P_2,$$

where P_2 is a Brauer-Severi surface bundle over $\sqrt[3]{(S, D)}$. Under ρ we have $\nu(\Sigma_3)$ contracting to a copy G in P_2 of the gerbe of the root stack, with $\mathcal{I}_G/\mathcal{I}_G^2$ locally free of rank 4. An exact sequence analogous to (5.2) shows that $\dim(\hat{\mathfrak{n}}/\hat{\mathfrak{n}}^2) = n + 3$, where $\hat{\mathfrak{n}}$ is the maximal ideal of the local ring, after base change to an affine étale chart of $\sqrt[3]{(S, D)}$, at a closed point \hat{r} over G ; in particular, P_2 has hypersurface singularities.

As above, we may work over $\text{Spec}(B)$ and perform previous analysis to show that 3 times each \mathbb{F}_1 component over the gerbe of the root stack is a Cartier divisor. This leads to an étale local defining equation at a point \hat{r} as above of $\delta_x\delta_y\delta_z - t^3$, with analogous notation to that appearing in (5.3). With notation as before, we have an affine chart $\text{Spec}(C)$ of

$$\hat{P}_2 := \text{Spec}(B) \times_S P_2,$$

which we may without loss of generality take to be μ_3 -invariant, and a closed immersion

$$\text{Spec}(C) \rightarrow \text{Spec}(\Delta)$$

identifying the local ring $C_{\hat{r}}$ with $\Delta_{\hat{\mathfrak{n}}'}/(\delta_x\delta_y\delta_z - t^3)$, where $\hat{\mathfrak{n}}'$ denotes the maximal ideal, corresponding to the image of \hat{r} in $\text{Spec}(\Delta)$. Although Δ is not determined canonically, its strict henselization at $\hat{\mathfrak{n}}'$ is canonically determined (up to fixing a separable closure of the residue field): it is a strict henselization of affine 3-space over B' .

By Proposition 2.5, there exists a scheme P_3 , Fano over S , with

$$\sqrt[3]{(S, D)} \times_S P_3 \cong P_2.$$

By Remark 2.6, $P_2 \rightarrow P_3$ is a coarse moduli space, so with the above notation, P_3 has affine coordinate ring C^{μ_3} . We claim, P_3 is regular; equivalently,

$$\hat{P}_3 := \text{Spec}(B) \times_S P_3$$

is regular. It suffices to verify the claim after passing to a strict henselization $(C^{\mu_3})^{sh}$ of P_3 . By [32, 18.8.10], we have

$$C \otimes_{C^{\mu_3}} (C^{\mu_3})^{sh} \cong C^{sh}.$$

Then a strict Henselization of P_2 (for Henselization of a stack, see [5, §2.7]) takes the form

$$[\mathrm{Spec}(C)/\mu_3] \times_{\mathrm{Spec}(C^{\mu_3})} \mathrm{Spec}((C^{\mu_3})^{sh}) \cong [\mathrm{Spec}(C^{sh})/\mu_3].$$

We have, μ_3 -equivariantly,

$$C^{sh} \cong \Delta^{sh}/(\delta_x \delta_y \delta_z - t^3)$$

at a point where three components meet. Hence upon taking μ_3 -invariants,

$$(C^{\mu_3})^{sh} \cong (C^{sh})^{\mu_3} \cong (\Delta^{sh})^{\mu_3}/(\delta_x \delta_y \delta_z - f).$$

The ring $(\Delta^{sh})^{\mu_3}$ is a strict Henselization of affine 3-space over B and hence is regular. A similar analysis takes care of the regularity at a point where two components meet. So, P_3 is regular.

6. REVERSE CONSTRUCTION

In this section, we start with a mildly degenerating Brauer-Severi surface bundle and carry out the reverse construction of Proposition 4.4.

For every $s \in D$, the fiber X_s has, geometrically, three irreducible components. Lemma 2.9 determines a degree 3 finite étale cover

$$\tilde{D} := \pi_0(X^{\mathrm{sm}}/S) \times_S D \rightarrow D.$$

For $s \in D$, let $\mathrm{Spec}(B) \rightarrow S$ be an étale neighborhood of s that trivializes $\tilde{D} \rightarrow D$. By Remark 2.10, the three sections of

$$\mathrm{Spec}(B) \times_S \tilde{D} \rightarrow \mathrm{Spec}(B) \times_S D$$

determine three divisors W_1, W_2, W_3 on $\mathrm{Spec}(B) \times_S X$, whose intersections with the fiber X_s of any $s \in \mathrm{Spec}(B) \times_S D$ are the irreducible components of X_s . Let w_1, w_2, w_3 be respective local defining equations of the divisors near a singular point $r \in X_s$. Then, after modification by a unit, we have $f = w_1 w_2 w_3$.

We claim, there is an étale local equation

$$w_1 w_2 w_3 - f, \quad \text{respectively,} \quad w_1 w_2 - f,$$

(after suitable renumbering of indices) for $\mathrm{Spec}(B) \times_S X$ in \mathbb{A}_B^3 , according to whether x is a point of X_s where 3 or 2 components meet.

To establish the claim, we argue as in Section 5. Let $\mathfrak{m} \subset B$ be the maximal ideal corresponding to s and $n = \dim(B_{\mathfrak{m}})$. We may choose a regular system of parameters for $B_{\mathfrak{m}}$ of the form f, g_1, \dots, g_{n-1} . The space $\mathfrak{n}/\mathfrak{n}^2$ has dimension $n+2$. Since the fiber X_s has hypersurface singularities, at a singular point $r \in X_s$ with corresponding maximal ideal $\mathfrak{n} \subset \mathcal{O}_{X,r}$ we have

$$\dim(\mathfrak{n}/(\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)) = 3.$$

The claim follows, as in Section 5, once we show that $\mathfrak{n}/(\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)$ is spanned by w_1, w_2, w_3 , respectively by two of them and a local parameter for the component \mathbb{P}^1 of the singular locus of the fiber.

We have

$$\dim((\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)/\mathfrak{n}^2) = n - 1.$$

The image of f in $\mathcal{O}_{X,r}$ lies in \mathfrak{n}^2 . Indeed, if not, then $\mathcal{O}_{X,r}/(f)$ would be regular, hence also the further localization, where we pass to the residue field at the generic point of D , would be regular. But the relative singular locus of $X \times_S D$ over D is flat (since it has constant Hilbert polynomial), hence some point of the singular locus over the generic point of D specializes to r , contradiction. So, the images of g_1, \dots, g_{n-1} form a basis of $(\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)/\mathfrak{n}^2$. Now, modulo g_1, \dots, g_{n-1} , we still have $f = w_1 w_2 w_3$, where $f = 0$ now defines the fiber X_s , and hence w_1, w_2, w_3 define its three irreducible components, each with multiplicity one.

Suppose, first, that r is the point where all three components meet. Then, for all $\{i, j, k\} = \{1, 2, 3\}$ the classes of w_j and w_k are a basis of $\mathfrak{n}/(\mathfrak{m}\mathcal{O}_{X,r} + w_i\mathcal{O}_{X,r} + \mathfrak{n}^2)$. So, the classes of w_1, w_2, w_3 are a basis of $\mathfrak{n}/(\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)$, and we are done in this case.

If r is a point where two components meet, say, those defined by w_1 and w_2 , then for $\{i, j\} = \{1, 2\}$ and local parameter p for the component \mathbb{P}^1 of the singular locus, w_j and p are a basis of $\mathfrak{n}/(\mathfrak{m}\mathcal{O}_{X,r} + w_i\mathcal{O}_{X,r} + \mathfrak{n}^2)$. It follows that w_1, w_2, p span $\mathfrak{n}/(\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)$, and we are done in this case as well.

The claim implies the corresponding singularity type of $\sqrt[3]{(S, D)} \times_S X$, where we have $f = t^3$ in the coordinate ring of an affine étale chart. The two blowing up steps therefore supply a resolution of singularities of $\sqrt[3]{(S, D)} \times_S X$. The three contractions then may be performed, as in Section 5, noticing by the normal cone description of Proposition A.9 that the outcome of each contraction is regular.

7. REGULAR DEGENERACY LOCUS IMPLIES MILD DEGENERATION

In this section we complete the proof of Proposition 4.4 by showing that a Brauer-Severi surface bundle $\pi: X \rightarrow S$ with X regular and singular fibers precisely along a regular divisor $D \subset S$, always has mild degeneration. We may assume that S is quasi-compact and integral. We consider various cases, depending on the dimension of S .

If $\dim(S) = 1$ then $\dim(X) = 3$, hence if $s \in S$ is any closed point then 3 is an upper bound on the embedding dimension of X_s at any of its closed points. So $X \rightarrow S$ has mild degeneration.

If $\dim(S) = 2$ then by the previous case we know that the fiber at the generic point of every component of D is, geometrically, a union of three

Hirzebruch surfaces. So, there is an open subscheme S' of S such that $S' \times_S X \rightarrow S'$ has mild degeneration and $\dim(S \setminus S') = 0$.

We apply the reverse construction (Section 6) to $S' \times_S X \rightarrow S'$ to obtain a smooth \mathbb{P}^2 -bundle

$$P' \rightarrow S' \times_S \sqrt[3]{(S, D)}.$$

By Proposition 2.1 (iii), this extends to a \mathbb{P}^2 -bundle

$$P \rightarrow \sqrt[3]{(S, D)}.$$

We now apply the forwards construction to obtain a mildly degenerating Brauer-Severi surface bundle $\hat{\pi}: \hat{X} \rightarrow S$.

By Remark 4.2 and the fact that a line bundle, ample on fibers, is relatively ample, we have the ampleness of the dual of the relative dualizing sheaf of $\hat{X} \rightarrow S$. This holds as well for $X \rightarrow S$, where the ampleness on geometric fibers over $S \setminus S'$ is dealt with by [30, 4.5.14], which tells us that ampleness of a line bundle on a Noetherian scheme is independent of any non-reduced structure, in combination with the fact that a cone over a twisted cubic curve has Picard group isomorphic to \mathbb{Z} .

Since

$$X \cong \text{Proj} \left(\bigoplus_{n \geq 0} \pi_*(\omega_{X/S}^\vee)^n \right) \quad \text{and} \quad \hat{X} \cong \text{Proj} \left(\bigoplus_{n \geq 0} \hat{\pi}_*(\omega_{\hat{X}/S}^\vee)^n \right),$$

where the direct image sheaves are locally free for all $n \gg 0$ with isomorphic restrictions over S' , they are isomorphic for all $n \gg 0$, hence $X \cong \hat{X}$ is a Brauer-Severi surface bundle with mild degeneration.

We observe that the argument for $\dim(S) = 2$ does not use the regularity of X , but only the flatness of $X \rightarrow S$ and the regularity of $S' \times_S X$.

The case $\dim(S) \geq 3$ reduces to the case $\dim(S) = 2$ by slicing the base. Let $s \in S$ be a point where the geometric fiber is irreducible with reduced subscheme a cone over a twisted cubic curve. Let f, g_1, \dots, g_{d-1} be a regular system of parameters for $\mathcal{O}_{S,s}$, and let T be the regular two-dimensional scheme $\text{Spec}(\mathcal{O}_{S,s}/(g_2, \dots, g_{d-1}))$. Let $Y := T \times_S X$. We claim, $Y \setminus Y_s$ is regular. This is clear at points in the relative smooth locus of $Y \setminus Y_s \rightarrow T \setminus \{s\}$. Consider the generic point η of $T \cap D$ with corresponding maximal ideal $\mathfrak{m} \subset \mathcal{O}_{S,\eta}$, and a closed point $r \in (X_\eta)^{\text{sing}}$, with corresponding maximal ideal $\mathfrak{n} \subset \mathcal{O}_{X,r}$. We use the exact sequence

$$0 \rightarrow (\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)/\mathfrak{n}^2 \rightarrow \mathfrak{n}/\mathfrak{n}^2 \rightarrow \mathfrak{n}/(\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2) \rightarrow 0.$$

The space in the middle has dimension $d+1$, on the right, 3, hence on the left, $d-2$. The space $\mathfrak{m}/\mathfrak{m}^2$ has dimension $d-1$ and is spanned by f, g_2, \dots, g_{d-1} . Moreover, f lies in the kernel of

$$\mathfrak{m}/\mathfrak{m}^2 \rightarrow (\mathfrak{m}\mathcal{O}_{X,r} + \mathfrak{n}^2)/\mathfrak{n}^2.$$

So g_2, \dots, g_{d-1} are linearly independent in $\mathfrak{n}/\mathfrak{n}^2$, and hence $\mathcal{O}_{Y,r}$ is regular.

By the observation made after the argument for the case $\dim(S) = 2$, we may conclude that $Y \rightarrow T$ has mild degeneration, and this is a contradiction.

8. LOCAL ANALYSIS I

For the next result we use the notation of Section 5 but work with schemes with μ_3 -action: P' will denote $\text{Proj}(B'[u, v, w])$, so $\widehat{P} = [P'/\mu_3]$, etc.

Proposition 8.1. *With $X \rightarrow S$ as in Proposition 4.4, the line bundle sheaf $\omega_{X/S}^\vee$ is relatively very ample over S , its direct image is locally free of rank 10, and formation of the direct image commutes with base change to an arbitrary Noetherian scheme over S . Furthermore, if we employ notation as remarked above and identify the the pullback of $\omega_{P'_2/\text{Spec}(B')}^\vee$ to $B\ell_{\tilde{P}'_\lambda}(B\ell_{P'_c}(B\ell_{P'_\mu} P'))$ with*

$$\omega^\vee - 2E_1 - E_2 - E_3$$

and sections with cubic forms in u, v, w vanishing appropriately along the E_i , then the sections form a free B' -module with basis

$$t^3w^3, \quad t^2uw^2, \quad tvw^2, \quad tu^2w, \quad uvw, \quad t^2v^2w, \quad t^3u^3, \quad t^2u^2v, \quad tuv^2, \quad t^3v^3.$$

The identification of line bundles in Proposition 8.1 comes from the observation that the line bundle associated with $\omega^\vee - 2E_1 - E_2 - E_3$ is the dual of the relative dualizing sheaf of $B\ell_{\tilde{P}'_\lambda}(B\ell_{P'_c}(B\ell_{P'_\mu} P'))$ over $\text{Spec}(B')$ (by standard behavior of dualizing sheaf under blowing up) and under the morphism to P'_2 has direct image $\omega_{P'_2/\text{Spec}(B')}^\vee$.

Proof. The observation from Remark 4.2, that the line bundle $\omega_{P'_2/\text{Spec}(B')}^\vee$ on fibers is very ample with space of global sections of dimension 10, implies the assertions in the first statement.

For the remaining assertion, we compute with local charts. We work, first, over the open subscheme

$$W' := \text{Spec}(B'[u, v])$$

of P' and use notation with the letter P replaced by W to denote restriction over W' . The blowup $B\ell_{W'_\mu} W'$ is covered by three charts:

Chart 1: coordinates u, t_1, v_1 with $t = ut_1$ and $v = uv_1$. Equation for E_1 : $u = 0$; equation for W'_c : $u = v_1 = 0$.

Chart 2: coordinates v, t_2, u_2 with $t = vt_2$ and $u = vu_2$. Equation for E_1 : $v = 0$; trivial intersection with W'_c , equation for \widetilde{W}'_λ : $t_2 = u_2 = 0$.

Chart 3: coordinates t, u_3, v_3 with $u = tu_3$ and $v = tv_3$. Equation for E_1 : $t = 0$; equation for W'_c : $t = v_3 = 0$.

The next blowup $B\ell_{W'_c}(B\ell_{W'_\mu} W')$ is covered by five charts, Chart 2 and:

Chart 1': coordinates u, t_1, v'_1 with $v_1 = uv'_1$. Equation for E_2 : $u = 0$; equation for \widetilde{W}'_λ : $t_1 = v'_1 = 0$.

Chart 1'': coordinates $t_1, v_1, \widetilde{u}''$ with $u = v_1u''$. Equation for E_2 : $v_1 = 0$; trivial intersection with \widetilde{W}'_λ .

Chart 3': (analogous)

Chart 3'': (analogous)

The next blowup $B\ell_{\widetilde{W}'_\lambda}(B\ell_{W'_c}(B\ell_{W'_\mu} W'))$ is covered by seven charts, of which we need just two:

Chart 1'_a: coordinates $u, t_1, v'_{1,a}$ with $v'_1 = t_1v'_{1,a}$. Equation for E_3 : $t_1 = 0$.

Chart 2_a: coordinates $v, t_2, u_{2,a}$ with $u_2 = t_2u_{2,a}$. Equation for E_3 : $t_2 = 0$

The union of charts 1'_a and 2_a contains the generic point of every component of E_1, E_2 , and E_3 over W' . With cyclic shifts of variables u, v, w we have additional affine opens U' and V' of P' with analogous charts of the blow-ups. By considering the analogous charts, too, we find that sections of $\omega^\vee - 2E_1 - E_2 - E_3$ are cubic forms in u, v, w with coefficients in B' , such that after applying any cyclic permutation of u, v, w and setting w to 1 the image in the coordinate ring of Chart 1'_a lies in the ideal

$$u^3t_1B[u, t_1, v'_{1,a}]/(u^3t_1^3 - f),$$

and the image in the coordinate ring of Chart 2_a lies in the ideal

$$v^2t_2B[v, t_2, u_{2,a}]/(v^3t_2^3 - f).$$

We compute with respect to the basis of degree 3 monomials in u, v, w . The kernel of

$$B'^{10} \rightarrow (B/f)[u, t_1, v'_{1,a}]/(u^3t_1)$$

may be computed with the help of the observation that the map factors through $((B/f)[t]/(t^3))^{10}$, which along with $(B/f)[u, t_1, v'_{1,a}]/(u^3t_1)$ is a free B/f -module. The kernel of

$$B'^{10} \rightarrow (B/f)[v, t_2, u_{2,a}]/(v^2t_2)$$

may be computed similarly. These computations lead to the determination of the module of such forms as the free module with the claimed basis. \square

9. LOCAL ANALYSIS II

We exhibit a Brauer-Severi surface fibration over a two-dimensional base, with singular fibers along a union of two intersecting divisors and smooth total space. For conic bundles this is easy:

$$\mathrm{Proj}(k[s, t, x, y, z]/(sx^2 + ty^2 - z^2)).$$

This conic bundle over $\mathbb{A}^2 = \mathrm{Spec}(k[s, t])$ has smooth total space and singular fibers along the divisor defined by $st = 0$.

For Brauer-Severi surface bundles this is more complicated. Over the generic point, the fiber is an anticanonically embedded Brauer-Severi surface, defined in \mathbb{P}^9 by

$$\frac{10 \cdot 11}{2} - h^0(\mathbb{P}^2, (\omega_{\mathbb{P}^2}^\vee)^2) = 55 - 28 = 27$$

quadratic equations. We will obtain equations for a Brauer-Severi surface bundle over \mathbb{A}^2 with smooth total space and singular fibers along the union of coordinate axes by applying results from §4. The existence of such a Brauer-Severi surface bundle has been shown by Maeda [55] using different methods (ideals and orders in a central simple algebra).

Let k be a field of characteristic different from 3 containing a primitive cube root of unity ζ , and let $\mu_3 \times \mu_3$ act on $\mathbb{A}^2 \times \mathbb{P}^2$ over k , where cube root of unity ζ in the first factor acts by

$$(s, t, u : v : w) \mapsto (\zeta s, t, \zeta u : \zeta^2 v : w)$$

and in the second factor acts by

$$(s, t, u : v : w) \mapsto (s, \zeta t, w : u : v).$$

Then

$$[(\mathbb{A}^2 \setminus \{0\}) \times \mathbb{P}^2 / \mu_3 \times \mu_3]$$

is a smooth \mathbb{P}^2 -fibration satisfying the hypotheses of Proposition 4.4. There is a corresponding mildly degenerating Brauer-Severi surface bundle

$$\pi: X \rightarrow \mathbb{A}^2 \setminus \{0\}. \tag{9.1}$$

By Proposition 8.1 applied to the pullback

$$\pi': X' \rightarrow \mathbb{A}^2 \setminus \{0\} \tag{9.2}$$

of π by $(s, t) \mapsto (s^3, t^3)$, first restricted over $\mathbb{A}^1 \times (\mathbb{A}^1 \setminus \{0\})$ and then restricted $(\mathbb{A}^1 \setminus \{0\}) \times \mathbb{A}^1$, the space of global sections of $\omega_{X'/\mathbb{A}^2 \setminus \{0\}}^\vee$ is the intersection of

$$\mathcal{S} :=$$

$$k[s, t, t^{-1}] \langle s^3w^3, s^2uw^2, svw^2, su^2w, uvw, s^2v^2w, s^3u^3, s^2u^2v, suv^2, s^3v^3 \rangle$$

and

$$\mathcal{T} :=$$

$$k[s, s^{-1}, t] \langle t^3 \tilde{w}^3, t^2 \tilde{u} \tilde{w}^2, t \tilde{v} \tilde{w}^2, t \tilde{u}^2 \tilde{w}, \tilde{u} \tilde{v} \tilde{w}, t^2 \tilde{v}^2 \tilde{w}, t^3 \tilde{u}^3, t^2 \tilde{u}^2 \tilde{v}, t \tilde{u} \tilde{v}^2, t^3 \tilde{v}^3 \rangle,$$

where

$$\begin{aligned}\tilde{u} &= \zeta u + \zeta^2 v + w, \\ \tilde{v} &= \zeta^2 u + \zeta v + w, \\ \tilde{w} &= u + v + w.\end{aligned}$$

Indeed, the action of cyclically permuting the coordinates u, v, w is compatible with the diagonal action by $\zeta, \zeta^2, 1$ on $\tilde{u}, \tilde{v}, \tilde{w}$.

Only nonnegative powers of s and t show up in an element of $\mathcal{S} \cap \mathcal{T}$. For a polynomial $f \in k[s, t, u, v, w]$, homogeneous of degree 3 in u, v, w :

- if $s^3 \mid f$, then $f \in \mathcal{S}$;
- if $t^3 \mid f$, then $f \in \mathcal{T}$;
- generally, if we write

$$f = \sum_{a,b \geq 0} s^a t^b f_{ab}, \quad \text{with } f_{ab} \in k[u, v, w],$$

for all a and b , then $f \in \mathcal{S}$ if and only if $s^a f_{ab} \in \mathcal{S}$ for all a and b , and $f \in \mathcal{T}$ if and only if $t^b f_{ab} \in \mathcal{T}$ for all a and b .

We easily characterize the homogeneous $f \in k[u, v, w]$ of degree 3 with $s^a f \in \mathcal{S}$ for $a \leq 2$:

a	k -basis
0	uvw
1	above and u^2w, vw^2, uv^2
2	above and uw^2, v^2w, u^2v

Similarly, with a bit of linear algebra we characterize the homogeneous $f \in k[u, v, w]$ of degree 3 with $t^b f \in \mathcal{T}$ for $b \leq 2$:

b	k -basis
0	$u^3 + v^3 + w^3 - 3uvw$
1	and $u^2w + \zeta vw^2 + \zeta^2 uv^2, uw^2 + \zeta v^2w + \zeta^2 u^2v, u^3 + \zeta^2 v^3 + \zeta w^3$
2	and $u^2w + \zeta^2 vw^2 + \zeta uv^2, uw^2 + \zeta^2 v^2w + \zeta u^2v, u^3 + \zeta v^3 + \zeta^2 w^3$

With these data, we determine the space of f with $s^a t^b f \in \mathcal{S} \cap \mathcal{T}$ for $0 \leq a, b \leq 2$ as trivial when $\min(a, b) = 0$ and of dimension $a + b - 1$ when $1 \leq a, b \leq 2$. E.g., for $a = b = 1$ the only linear combination of uvw, u^2w, vw^2, uv^2 that belongs to the space indicated for $b = 1$ is, up

to scalar multiplication, $u^2w + \zeta vw^2 + \zeta^2 uv^2$. We deduce that $\mathcal{S} \cap \mathcal{T}$ is the free $k[s, t]$ -module with basis

$$\begin{aligned} & t^3uvw, \quad st(u^2w + \zeta vw^2 + \zeta^2 uv^2), \quad st^2(u^2w + \zeta^2 vw^2 + \zeta uv^2), \\ & st^3(u^2w + vw^2 + uv^2), \quad s^2t(uw^2 + \zeta v^2w + \zeta^2 u^2v), \\ & s^2t^2(uw^2 + \zeta^2 v^2w + \zeta u^2v), \quad s^2t^3(uw^2 + v^2w + u^2v), \\ & s^3(u^3 + v^3 + w^3 - 3uvw), \quad s^3t(u^3 + \zeta^2 v^3 + \zeta w^3), \quad s^3t^2(u^3 + \zeta v^3 + \zeta^2 w^3). \end{aligned}$$

By Proposition 8.1, the basis of $\mathcal{S} \cap \mathcal{T}$ defines a morphism $X' \rightarrow \mathbb{A}^2 \times \mathbb{P}^9$. With linear algebra, we find $f_1, \dots, f_{27} \in k[s, t, x_0, \dots, x_9]$ that are

- homogeneous of degree 2 in x_0, \dots, x_9 and at most linear in s, t .
- zero upon substituting s^3 for s , t^3 for t , and the i th basis element for x_{i-1} , for every i ,
- linearly independent over $k(s, t)$.

The polynomials are displayed in Table 1. The conditions imply that

$$\text{Proj}(k(s, t)[x_0, \dots, x_9]/(f_1, \dots, f_{27}))$$

is the Brauer-Severi surface that appears as generic fiber of π .

Lemma 9.1. *Let f_1, \dots, f_{27} be as in Table 1, where k is a field of characteristic different from 3 with primitive cube root of unity ζ . Then $k[s, t, x_0, \dots, x_9]/(f_1, \dots, f_{27})$ is finitely generated and free as a module over $k[s, t, x_0, x_1, x_7]$, with basis 1, $x_2, x_3, x_4, x_5, x_5^2, x_6, x_8, x_9$.*

Proof. Using the relations $f_i = 0$ for $i = 1, \dots, 27$ we write down 9×9 -matrices for the action of multiplication by x_i on the claimed basis elements for $i \in \{2, 3, 4, 5, 6, 8, 9\}$. The matrices, as may be checked, commute pairwise and obey relations corresponding to f_1, \dots, f_{27} . So the evident homomorphism from $k[s, t, x_0, \dots, x_9]/(f_1, \dots, f_{27})$ to the free $k[s, t, x_0, x_1, x_7]$ -module of rank 9 is an isomorphism. \square

Lemma 9.1 implies that the ideal (f_1, \dots, f_{27}) of $k[s, t, x_0, \dots, x_9]$ is prime. Indeed, the ideal is saturated with respect to x_0, \dots, x_9 (e.g., $x_0g \in (f_1, \dots, f_{27})$ implies $g \in (f_1, \dots, f_{27})$ for $g \in k[s, t, x_0, \dots, x_9]$), and the ideal generated by f_1, \dots, f_{27} in $k(s, t)[x_0, \dots, x_9]$ defines an anticanonically embedded Brauer-Severi variety. So the latter ideal, and hence also the former ideal, is prime.

Proposition 9.2. *Let k be a field of characteristic different from 3 with primitive cube root of unity ζ , $\pi: X \rightarrow \mathbb{A}^2 \setminus \{0\}$ and $\pi': X' \rightarrow \mathbb{A}^2 \setminus \{0\}$ as in (9.1)–(9.2) and $f_1, \dots, f_{27} \in k[s, t, x_0, \dots, x_9]$ as in Table 1. The coherent sheaf $\pi_*\omega_\pi^\vee$ is free of rank 10, and for some basis the scheme-theoretic image in $\mathbb{A}^2 \times \mathbb{P}^9$ is defined by the prime ideal (f_1, \dots, f_{27}) and is a nonsingular variety, flat over \mathbb{A}^2 . For an explicit basis of the free*

i	f_i
1	$-3\zeta^2x_0x_4 - x_1x_3 + x_2^2$
2	$-tx_1^2 + 3\zeta x_0x_5 + x_2x_3$
3	$-tx_1x_2 - 3x_0x_6 + x_3^2$
4	$(\zeta - \zeta^2)x_0x_7 - x_1x_5 + x_2x_4$
5	$(1 - \zeta^2)x_0x_8 - x_1x_6 + x_3x_4$
6	$-(\zeta - \zeta^2)x_0x_9 - tx_1x_4 + x_3x_5$
7	$\zeta^2x_1x_6 + \zeta x_3x_4 + x_2x_5$
8	$\frac{1-\zeta^2}{3}x_1x_8 - \frac{1-\zeta^2}{3}x_2x_7 + x_4^2$
9	$\frac{1-\zeta}{3}x_1x_9 - \frac{1-\zeta}{3}x_3x_7 + x_4x_5$
10	$\zeta^2tx_1x_4 + \zeta x_3x_5 + x_2x_6$
11	$-x_3x_7 + (1 - \zeta)x_4x_5 + x_2x_8$
12	$-tx_1x_7 + (1 - \zeta^2)x_5^2 + x_2x_9$
13	$-9sx_0^2 + \zeta tx_1x_5 + \zeta^2tx_2x_4 + x_3x_6$
14	$-3(1 - \zeta^2)sx_0x_1 - tx_1x_7 + (1 - \zeta)x_5^2 + x_3x_8$
15	$-3(1 - \zeta)sx_0x_2 - tx_2x_7 + (1 - \zeta^2)tx_4^2 + x_3x_9$
16	$3\zeta sx_0x_1 - x_5^2 + x_4x_6$
17	$-(1 - \zeta)sx_1^2 - \zeta x_5x_7 + x_4x_8$
18	$-(1 - \zeta^2)sx_1x_2 - \zeta^2x_6x_7 + x_4x_9$
19	$3\zeta^2sx_0x_2 - tx_4^2 + x_5x_6$
20	$-(1 - \zeta)sx_1x_2 - \zeta x_6x_7 + x_5x_8$
21	$-(1 - \zeta^2)sx_2^2 - \zeta^2tx_4x_7 + x_5x_9$
22	$-3sx_0x_3 - tx_4x_5 + x_6^2$
23	$-(1 - \zeta^2)sx_1x_3 + (\zeta - \zeta^2)sx_2^2 - \zeta tx_4x_7 + x_6x_8$
24	$-(1 - \zeta)sx_2x_3 - \zeta tx_4x_8 + x_6x_9$
25	$-3\zeta sx_1x_4 - x_7x_9 + x_8^2$
26	$-3\zeta^2sx_1x_5 - 3\zeta sx_2x_4 - tx_7^2 + x_8x_9$
27	$-3\zeta^2sx_2x_5 - tx_7x_8 + x_9^2$

TABLE 1. The polynomials $f_1, \dots, f_{27} \in k[s, t, x_0, \dots, x_9]$.

coherent sheaf $\pi'_*\omega_{\pi'}^\vee$ of rank 10 the scheme-theoretic image in $\mathbb{A}^2 \times \mathbb{P}^9$ is defined by the ideal obtained from (f_1, \dots, f_{27}) by applying $s \mapsto s^3$ and $t \mapsto t^3$, which is also prime. The fiber over $s = t = 0$ is an irreducible, nonreduced scheme whose reduced subscheme is a cone over a twisted cubic curve.

The explicit basis mentioned in the statement of Proposition 9.2 is the one given above, as the basis of $\mathcal{S} \cap \mathcal{T}$.

Proof. As remarked after Lemma 9.1, the ideal generated by f_1, \dots, f_{27} is prime, and the same holds after applying $s \mapsto s^3$ and $t \mapsto t^3$. These

facts, plus the remark about $\text{Proj}(k(s,t)[x_0, \dots, x_9]/(f_1, \dots, f_{27}))$ made before Lemma 9.1, justify the statements about scheme-theoretic image.

We directly analyze the fiber over $s = t = 0$. First, the reduced subscheme is contained in the locus $x_3 = x_5 = x_6 = x_8 = x_9 = 0$, as we see directly from the defining equations. The defining equations with

$$s = t = x_3 = x_5 = x_6 = x_8 = x_9 = 0$$

no longer involve x_1 . There are three equations, quadratic in x_0, x_2, x_4, x_7 , the equations of a twisted cubic. So the reduced subscheme of the fiber is a cone over a twisted cubic. The coordinate point

$$p_0 = (1 : 0 : 0 : 0 : 0 : 0 : 0 : 0 : 0 : 0)$$

lies on the fiber and has Zariski tangent space of dimension 3. But p_0 is smooth point of the reduced subscheme. So, the fiber is nonreduced.

Now we consider the variety

$$\text{Proj}(k[s, t, x_0, \dots, x_9]/(f_1, \dots, f_{27})).$$

Flatness over \mathbb{A}^2 follows from Lemma 9.1. It remains to show that this is a nonsingular variety. For this, we let the two-dimensional torus \mathbb{G}_m^2 act on \mathbb{A}^2 by $(\delta, \varepsilon) \cdot (s, t) = (\delta^3 s, \varepsilon^3 t)$ and on \mathbb{P}^9 by sending $(x_0 : \dots : x_9)$ to

$$(\varepsilon^3 x_0 : \delta \varepsilon x_1 : \delta \varepsilon^2 x_2 : \delta \varepsilon^3 x_3 : \delta^2 \varepsilon x_4 : \delta^2 \varepsilon^2 x_5 : \delta^2 \varepsilon^3 x_6 : \delta^3 x_7 : \delta^3 \varepsilon x_8 : \delta^3 \varepsilon^2 x_9)$$

and thereby obtain a torus action on the variety defined by f_1, \dots, f_{27} . Suppose the variety is singular. Since the singular locus is closed and torus-invariant, it must contain a fixed point, i.e., a coordinate point, of the fiber over $s = t = 0$. From the description above, we only have to consider the coordinate points p_0, p_1 , and p_7 . For each we specify 7 defining equations and 7 variables:

point	equations	variables
p_0	$f_1, f_2, f_3, f_4, f_5, f_6, f_{13}$	$s, x_4, x_5, x_6, x_7, x_8, x_9$
p_1	$f_1, f_2, f_4, f_5, f_8, f_9, f_{17}$	$s, t, x_3, x_5, x_6, x_8, x_9$
p_7	$f_4, f_8, f_9, f_{17}, f_{18}, f_{25}, f_{26}$	$t, x_0, x_2, x_3, x_5, x_6, x_9$

By direct computation we see that the corresponding Jacobian matrix at each p_i is nonsingular, and thus p_i is a nonsingular point of the variety defined by f_1, \dots, f_{27} . \square

The proof of Theorem 1.2 proceeds as in the case of conic bundles [64], where we have incorporated a modification suggested by D. Rydh that weakens the requirement of resolution of singularities (functoriality is not needed) and eliminates the need for destackification [16]:

- Represent the generic fiber of π by a 3-torsion Brauer group element $\alpha \in \text{Br}(k(S))$. Resolving the defining ideal of a closed subscheme, complement of some dense open to which α extends, we may suppose that α is ramified along a simple normal crossing divisor $D_1 \cup \dots \cup D_\ell$. Let $\mathcal{R} := \sqrt[3]{(S, \{D_1, \dots, D_\ell\})}$ be the iterated root stack; then by Remark 2.4, α is the restriction of an element

$$\beta \in \text{Br}(\mathcal{R})[3].$$

Pass to a μ_3 -gerbe $\mathcal{G} \rightarrow \mathcal{R}$ whose class is a lift $\beta_0 \in H^2(\mathcal{R}, \mu_3)$ of β . The restriction of π to $\pi^{-1}(U)$ is a smooth \mathbb{P}^2 -bundle by [40, Lem. 1.7]. So, there is a corresponding sheaf of Azumaya algebras over U and a locally free sheaf of rank 3 over $\mathcal{G} \times_{\mathcal{R}} U$. This spreads out to a coherent sheaf \mathcal{E} on \mathcal{G} . It is known [26, Lem. 1] that by blowing up an appropriate Fitting ideal sheaf of \mathcal{E} whose support is the locus where \mathcal{E} fails to be locally free, we obtain a locally free coherent sheaf on the blow-up whose restriction over $\mathcal{G} \times_{\mathcal{R}} U$ agrees with \mathcal{E} . The Fitting ideal sheaf descends to an ideal sheaf on \mathcal{R} , hence there is birational projective $\mathcal{R}' \rightarrow \mathcal{R}$, restricting to an isomorphism over U , and a locally free sheaf on $\mathcal{G} \times_{\mathcal{R}} \mathcal{R}'$ whose restriction over $\mathcal{G} \times_{\mathcal{R}} U$ agrees with \mathcal{E} .

- Apply Raynaud-Gruson flatification [66, §5] to $\mathcal{R}' \rightarrow S$, or better, to $T_\alpha \rightarrow S_\alpha$ for a Zariski covering (S_α) of S with $\mathcal{R}' \times_S S_\alpha \cong [T_\alpha / (\mu_3)^{n_\alpha}]$, to obtain a coherent sheaf of ideals \mathcal{I} defining a closed subscheme of S disjoint from U , such that for any proper birational $\psi: \tilde{S} \rightarrow S$ restricting to an isomorphism over U such that $\psi^{-1}\mathcal{I} \cdot \mathcal{O}_{\tilde{S}}$ is an invertible sheaf of ideals, the operation of scheme-theoretic closure on the level of atlases defines a closed substack of $\mathcal{R}' \times_S \tilde{S}$ that is flat over \tilde{S} . (All that we will use is that the substack is proper and quasi-finite over \tilde{S} .)
- Resolve the marked ideal consisting of \mathcal{I} and D_1, \dots, D_ℓ by some $\psi: \tilde{S} \rightarrow S$, composition of blow-ups along centers that are smooth and disjoint from U , and have normal crossings with D_1, \dots, D_ℓ (or their proper transforms along with the exceptional divisors of earlier blow-ups), so that $\psi^{-1}\mathcal{I} \cdot \mathcal{O}_{\tilde{S}}$ is an invertible sheaf of ideals. As a consequence of the universal property of blowing up, we see that there is a locally free coherent sheaf on the normalization of

$$\mathcal{G} \times_S \tilde{S}$$

whose restriction over $\mathcal{G} \times_{\mathcal{R}} U$ agrees with \mathcal{E} . The same holds if we replace the indicated fiber product by the base-change of \mathcal{G} to

the iterated root stack of \tilde{S} along the proper transforms of the divisors D_1, \dots, D_ℓ and the exceptional divisors of the blow-ups.

- Thus we may suppose that the coherent sheaf \mathcal{E} on \mathcal{G} is locally free and \mathcal{R} is an iterated root stack $\sqrt[3]{(\tilde{S}, \{\tilde{D}_1, \dots, \tilde{D}_m\})}$ over a smooth proper variety \tilde{S} . Now \mathcal{E} determines (cf. §2.2) a Brauer-Severi fibration over \mathcal{R} , which by abuse of notation we write as $\mathbb{P}(\mathcal{E}) \rightarrow \mathcal{R}$. We may suppose, for every i , that over a general point of \tilde{D}_i the stabilizer action on the fiber of $\mathbb{P}(\mathcal{E}) \rightarrow \mathcal{R}$ is balanced. If not, then the restriction of \mathcal{E} to

$$\mathcal{G} \times_{\mathcal{R}} \tilde{\mathfrak{D}}_i \quad (9.3)$$

splits canonically according to multiplicity as $\mathcal{E}_1 \oplus \mathcal{E}_2$ and, letting h denote the inclusion of (9.3) in \mathcal{G} , for $j = 1, 2$ the kernel $\tilde{\mathcal{E}}^{(j)}$ of

$$\mathcal{E} \rightarrow h_* \mathcal{E}_j$$

is locally free, with trivial stabilizer action on $\mathbb{P}(\tilde{\mathcal{E}}^{(j)})$ for suitable j over a general point of \tilde{D}_i . This contradicts the assumption that α is ramified along \tilde{D}_i .

- After blowing up intersections of pairs of rooted divisors, following the convention in §2.4 we arrive at a situation where no triple of divisors intersects, and for every pair of intersecting divisors, the corresponding projective representation $\mu_3 \times \mu_3 \rightarrow PGL_3$ over a general point of the intersection is faithful. The group-theoretic input is the fact that no more than two independent commuting copies of μ_3 may be found in PGL_3 , so any homomorphism $\mu_3 \times \mu_3 \times \mu_3 \rightarrow PGL_3$ has nontrivial kernel. We arrive at a configuration as claimed by blowing up intersections of pairs, first among triples for which the projective representation associated with every pair is faithful, then pairs with non-faithful projective representation.
- Apply Proposition 4.4 to obtain a mildly degenerating Brauer-Severi surface bundle with smooth total space

$$\pi_0: \tilde{X}_0 \rightarrow \tilde{S}_0, \quad (9.4)$$

where \tilde{S}_0 denotes the complement in \tilde{S} of $\bigcup_{i < j} \tilde{D}_i \cap \tilde{D}_j$. Proposition 8.1 tells us that $\mathcal{F}_0 := (\pi_0)_* \omega_{\pi_0}^\vee$ is locally free of rank 10, and its local sections embed \tilde{X}_0 in $\mathbb{P}(\mathcal{F}_0^\vee)$. By Lemma 2.8, over a suitable étale neighborhood of a point of $\tilde{D}_i \cap \tilde{D}_j$ ($i < j$) we may identify (9.4) with the product with $\mathbb{A}^{\dim(S)-2}$ of (9.1). It follows by Proposition 9.2 that, letting ι denote the inclusion $\tilde{S}_0 \rightarrow \tilde{S}$, the

coherent sheaf $\mathcal{F} := \iota_* \mathcal{F}_0$ (coherent since ι is the inclusion of the complement of a closed subset of codimension 2 [32, 5.11.1]) is in fact locally free, and the closure \tilde{X} of \tilde{X}_0 in $\mathbb{P}(\mathcal{F}^\vee)$ is a standard Brauer-Severi surface bundle.

10. APPLICATION TO RATIONALITY

Here we prove Theorem 1.4. We use a theorem of Voisin [75, Thm. 2.1], in the form of [23, Thm. 2.3]:

Theorem 10.1. *Let k be an uncountable algebraically closed field, B a variety over k , and $\phi: \mathcal{X} \rightarrow B$ a flat projective morphism with integral fibers and smooth generic fiber. Suppose that there exists a k -point $b_0 \in B$ such that the fiber $X_0 := \phi^{-1}(b_0)$ satisfies:*

- (O) *for some positive integer n , invertible in k , there is nontrivial unramified cohomology of μ_n on X_0 ;*
- (R) *X_0 admits a universally CH_0 -trivial resolution of singularities.*

Then, for every general $b \in B$, the fiber $X_b := \phi^{-1}(b)$ is not stably rational.

The reader may turn to [23] for the definitions of *universally CH_0 -trivial scheme* and *universally CH_0 -trivial morphism*. For instance, (R) is satisfied if X_0 admits a resolution of singularities $\tilde{X}_0 \rightarrow X_0$ which is a projective birational morphism, isomorphism over the complement of a curve, restricting to a generically smooth quadric surface bundle over the curve; or a composite of such morphisms.

For a proper geometrically integral scheme X over a field ℓ , say of dimension d , a related notion (equivalent, when X is smooth [23, Prop. 1.4]), is for X to admit a *Chow decomposition of the diagonal*. This means that in the Chow group $\text{CH}_d(X \times X)$ of d -dimensional cycles modulo rational equivalence, the class $[\Delta_X]$ of the diagonal of X is equal to a sum

$$Z + [X] \times z_0$$

for some d -dimensional cycle Z supported on $D \times X$, for a closed subset $D \subsetneq X$, and 0-cycle z_0 of degree 1 on X . If we let $\delta_X \in \text{CH}_0(X_{\ell(X)})$ denote the restriction of Δ_X to $X_{\ell(X)} := \text{Spec}(\ell(X)) \times_{\text{Spec}(\ell)} X$, this is equivalent to:

$$(*) \quad X \text{ has a 0-cycle } z_0 \text{ of degree 1 and } [\delta_X] = z_0 \text{ in } \text{CH}_0(X_{\ell(X)}).$$

Condition (R), with CH_0 -trivial resolution of singularities $\tilde{X}_0 \rightarrow X_0$, implies the equivalence of $(*)$ on X_0 and \tilde{X}_0 , hence the equivalence of admitting a Chow decomposition of the diagonal. Chow decomposition of the diagonal is obstructed by Condition (O) but implied by stable rationality, while an argument with Chow schemes [23, App. B] (Chow

functors in case $\text{char}(k) > 0$ [36, §3]) shows that the k -points whose fibers admit a Chow decomposition of the diagonal occupy a countable union of closed subsets of B . An alternative approach, bypassing Chow groups, appears in recent work [63], [44].

We return to Theorem 1.4, with $\text{char}(k) \neq 2, 3$, and smooth degree 2 del Pezzo surface S .

Remark 10.2. The ingredients from this paper, needed for the proof of Theorem 1.4 are:

- Theorem 1.2, existence of standard Brauer-Severi surface bundle models, albeit only in the case $\dim(S) = 2$ and 3-torsion element of $\text{Br}(k(S))$ ramified along a nonsingular divisor; special for the case of dimension 2 is the result on extension of a generically defined sheaf of Azumaya algebras, Proposition 2.1 (ii), and the conclusion of Theorem 1.2 is valid in this case with $\tilde{S} = S$.
- Proposition 4.4, dictionary between mildly degenerating Brauer-Severi surface bundles and smooth \mathbb{P}^2 -fibrations over associated root stacks.

We first treat the case $d = 3$ and then explain the modifications to deal with general d . The anticanonical morphism expresses S as double cover of \mathbb{P}^2 branched along a nonsingular quartic curve R , which we take to have defining equation h with respect to some coordinates.

Step 1. There exists a nonsingular cubic curve D_0 meeting R in 6 points of multiplicity 2, such that the 6 points lie on a conic. Among the 28 bitangent lines to R , at most 12 are hyperflexes (lines meeting R in a point of multiplicity 4), since the weighted count of Weierstrass points of R is 24 and each hyperflex contributes 2; cf. [73]. Bitangent lines are identified with odd theta characteristics, and it is known classically (see, e.g., [29, §2]) that any set of 8 bitangent lines, which we may suppose disjoint from the set of hyperflexes, contains a syzygetic triple ℓ, ℓ', ℓ'' : the six points of intersection of $\ell \cup \ell' \cup \ell''$ with R lie on a conic $f = 0$. Using the same symbols for defining linear forms and suitably rescaling h , we obtain by Max Noether's theorem [28, §5.5] a relation $h = f^2 + \ell\ell'\ell''\ell'''$ for a fourth bitangent ℓ''' . Let j be a line in general position with respect to ℓ, ℓ', ℓ'' with suitably scaled defining equation. Then we easily see that the cubic $g := \ell\ell'\ell'' - 2fj - j^2\ell'''$, appearing in the relation $h = (f + jl''')^2 + g\ell'''$, is nonsingular and meets R in six points of multiplicity 2.

Step 2. Let $\tilde{D}_0 \rightarrow D_0$ be a nontrivial cyclic degree 3 étale cover, determining (Remark 2.3) a 3-torsion element of $\text{Br}(\mathbb{P}^2 \setminus D_0)$, which extends (Remark 2.4) to an element of $\text{Br}(\sqrt[3]{(\mathbb{P}^2, D_0)})$, represented ([8],

[24]) by an index 3 division algebra at the generic point. This spreads out, as mentioned in Remark 10.2, to a sheaf of Azumaya algebras over $\sqrt[3]{(\mathbb{P}^2, D_0)}$. We obtain an element $\gamma_0 \in H^2(\sqrt[3]{(\mathbb{P}^2, D_0)}, \mu_3)$, gerbe $G_0 \rightarrow \sqrt[3]{(\mathbb{P}^2, D_0)}$ with this class, and locally free coherent sheaf E_0 of rank 3 on G_0 such that the pullback of the sheaf of Azumaya algebras is $\text{End}(E_0)$. We perform base change via $S \rightarrow \mathbb{P}^2$ to obtain $D = S \times_{\mathbb{P}^2} D_0$, a *reducible* curve $D = D_1 \cup D_2$ with étale degree 3 cover $\tilde{D} \rightarrow D$, element $\gamma \in H^2(\sqrt[3]{(S, D)}, \mu_3)$, gerbe $G \rightarrow \sqrt[3]{(S, D)}$, and locally free coherent sheaf E on G .

Step 3. Apply the deformation-theoretic result [36, Prop. 17] to exhibit an exact sequence $0 \rightarrow \tilde{E} \rightarrow E \rightarrow Q \rightarrow 0$, where Q is supported on a divisor and is locally free there of rank 2, such that the space of obstructions

$$\ker(H^2(\sqrt[3]{(S, D)}, \tilde{E}^\vee \otimes \tilde{E}) \rightarrow H^2(\sqrt[3]{(S, D)}, \mathcal{O}_{\sqrt[3]{(S, D)}}))$$

vanishes, where we abuse notation, as in loc. cit., and let $\tilde{E}^\vee \otimes \tilde{E}$ denote the corresponding sheaf of Azumaya algebras on $\sqrt[3]{(S, D)}$.

Step 4. Deduce as in [36, Sect. 5] that, taking b_0 to be the point corresponding to divisor D with $\tilde{D} \rightarrow D$ in the space B of reduced nodal curves on S with cyclic degree 3 étale cover we have:

- the corresponding divisor $\mathcal{D} \subset B \times S$, which is smooth,
- the root stack $\sqrt[3]{(B \times S, \mathcal{D})}$,
- upon replacing B by a suitable étale neighborhood of b_0 , a class

$$\Gamma \in H^2(\sqrt[3]{(B \times S, \mathcal{D})}, \mu_3)$$

restricting to γ ,

- a corresponding gerbe $\mathcal{G} \rightarrow \sqrt[3]{(B \times S, \mathcal{D})}$ restricting to G ,
- upon again passing to an étale neighborhood of b_0 , a locally free coherent sheaf $\tilde{\mathcal{E}}$ on \mathcal{G} restricting to \tilde{E} ,
- a smooth \mathbb{P}^2 -bundle $P \rightarrow \sqrt[3]{(B \times S, \mathcal{D})}$, which upon base change to \mathcal{G} yields $\mathbb{P}(\tilde{\mathcal{E}})$.

Step 5. Apply Proposition 4.4 to P to obtain a Brauer-Severi bundle $\pi: \mathcal{X} \rightarrow B \times S$ satisfying the hypotheses of Theorem 10.1. Since, by construction, the discriminant curve of the Brauer-Severi bundle $X_0 \rightarrow \{b_0\} \times S$ has two components and the Brauer class is given by cyclic covers, nontrivial on each component and étale over the nodes, condition (O) is satisfied: the unramified H^2 of μ_3 is the 3-torsion in the unramified Brauer group of X_0 , which is nontrivial since the elements $\gamma_i \in \text{Br}(k(S))[3]$ ramified along just D_i by the covering $\tilde{D}_0 \rightarrow D_0 \cong D_i$

(Remark 2.3) for $i = 1, 2$, satisfy: their sum pulls back to 0 in $\text{Br}(k(X_0))$, and each individually pulls back to a nontrivial unramified class. (See [22, §2] for a similar argument involving 2-torsion Brauer classes.) Condition (R) holds, since the singularities of X_0 consist of three curves over each node of the discriminant curve with the étale local form of the hypersurface singularity $uv = xyz$; a resolution is given by blowing up first one curve, then the proper transforms of the other two over each node, and each blow-up yields a morphism restricting to a generically smooth quadric surface bundle over a curve. Since $| -3K_S |$ contains curves with E_6 -singularity, we obtain as in the proof of [35, Lemma 3.1] (where the extra assumption involving a pair of curves meeting in an odd number of points is used only for the monodromy analysis mod 2 and is not needed here) that the space of nodal curves with étale cyclic degree 3 cover is irreducible and the proof of Theorem 1.4 is complete. (It is really for ease of argument that we use such a particular curve $D \in | -3K_S |$; a more involved argument where D can be an arbitrary union of two smooth curves of positive genus meeting transversely is given in [47].)

The case $d > 3$ is dealt with by taking D_3 to be the pre-image in S of a general curve $C \subset \mathbb{P}^2$ of degree $d - 3$ and defining $D'_0 := D_0 \cup C$. We pull back $\gamma_0 \in H^2(\sqrt[3]{(\mathbb{P}^2, D_0)}, \mu_3)$ and the gerbe G_0 to obtain $\gamma'_0 \in H^2(\sqrt[3]{(\mathbb{P}^2, D'_0)}^{\text{sm}}, \mu_3)$ and a gerbe G'_0 . The pull-back of the locally free coherent sheaf E_0 has trivial μ_3 -stabilizer actions over points of C . By performing suitable elementary transformations over the component of the gerbe of the root stack over C , we may obtain balanced actions, compatible with those over D_0 . Then the smooth \mathbb{P}^2 -bundle corresponding to the locally free sheaf has the same étale local isomorphism type as that corresponding to E (Step 2). We thereby obtain an extension of γ'_0 to $\gamma''_0 \in H^2(\sqrt[3]{(\mathbb{P}^2, D'_0)}, \mu_3)$, of G'_0 to $G''_0 \rightarrow \sqrt[3]{(\mathbb{P}^2, D'_0)}$, and of E to a locally free coherent sheaf on G''_0 . We proceed with Steps 3 and 4, taking D to be the union of D_1, D_2 , and the pull-back D_3 of C ; the triviality of the cyclic cover over D_3 poses no problems in carrying out these steps.

APPENDIX A. BIRATIONAL CONTRACTIONS

Let k be a field and X a scheme, proper over k . A line bundle L on X is called *semiample* if L^m is globally generated for some $m > 0$. Then the set $M(X, L)$ of $m \geq 0$, such that L^m is globally generated, consists, for $m \gg 0$, of precisely the multiples of a well-defined positive integer, the *exponent* of $M(X, L)$; cf. [52, §2.1.B]. We start by recalling the contraction determined by a semiample line bundle.

Proposition A.1. *Let X be a scheme, proper over a field k , with semi-ample line bundle L and sections $s_0, \dots, s_M \in H^0(X, L^m)$ generating L^m for some $m > 0$, thereby inducing a morphism $\psi: X \rightarrow \mathbb{P}^M$. Let $Y = \psi(X)$, and write the Stein factorization of ψ as morphism*

$$\pi: X \rightarrow X'$$

followed by finite morphism $f: X' = \text{Spec}(\psi_ \mathcal{O}_X) \rightarrow Y$. We let $\mathcal{O}_Y(1)$, respectively $\mathcal{O}_{X'}(1)$ denote the restriction to Y , respectively the pull-back to X' of the line bundle $\mathcal{O}_{\mathbb{P}^M}(1)$, so $\pi^* \mathcal{O}_{X'}(1) \cong L^m$. Then:*

- (i) *The morphism $\pi: X \rightarrow X'$ is proper and induces an isomorphism $\mathcal{O}_{X'} \xrightarrow{\sim} \pi_* \mathcal{O}_X$.*
- (ii) *The line bundle $\mathcal{O}_{X'}(1)$ is ample, and for any positive integer n such that $\mathcal{O}_{X'}(n)$ is very ample, the morphism $X \rightarrow \mathbb{P}^N$ determined by a basis of $H^0(X, L^{mn})$ factors uniquely through X' , identifying X' with the image of X in \mathbb{P}^N .*
- (iii) *With the property in (ii) the scheme X' , with $\pi: X \rightarrow X'$, is determined uniquely up to canonical isomorphism, independently of the choice of m and the generating sections, and the powers of L that are isomorphic to the pull-back of a line bundle from X' are precisely all multiples of the exponent of $M(X, L)$.*

Proof. We have stated, in (i), a basic property of the Stein factorization; see [31, 4.3.1]. The first statement in (ii) follows from standard facts on ample line bundles, and for the second statement, using the isomorphism $\mathcal{O}_{X'} \cong \pi_* \mathcal{O}_X$ we have

$$H^0(X, L^{mn}) \cong H^0(X', \mathcal{O}_{X'}(n)), \quad (\text{A.1})$$

so for n such that $\mathcal{O}_{X'}(n)$ is very ample, a choice of basis compatibly maps X to \mathbb{P}^N and embeds X' in \mathbb{P}^N . The factorization is unique, and by (i), identifies X' with the image of $X \rightarrow \mathbb{P}^N$. For (iii), if $\tilde{s}_0, \dots, \tilde{s}_{\tilde{M}}$ is another collection of generating sections of L^m , leading to $\tilde{\pi}: X \rightarrow \tilde{X}'$, then for n such that $\mathcal{O}_{X'}(n)$ and $\mathcal{O}_{\tilde{X}'}(n)$ are very ample, X' and \tilde{X}' are both identified with the image of $X \rightarrow \mathbb{P}^N$, as in (ii). The isomorphism $X' \cong \tilde{X}'$ obtained this way is independent of n : it suffices to compare the isomorphisms arising from n , leading to $X \rightarrow \mathbb{P}^N$, and dn , for some $d > 0$, leading to $X \rightarrow \mathbb{P}^{N'}$, but these are compatible with the d -tuple Veronese embedding of \mathbb{P}^N and the rational map from $\mathbb{P}^{N'}$ corresponding to

$$\text{Sym}^d(H^0(X, L^{mn})) \rightarrow H^0(X, L^{dmn}).$$

When $\pi: X \rightarrow X'$ and $\tilde{\pi}: X \rightarrow \tilde{X}'$ arise from different powers of L the argument is similar, with pairs n and \tilde{n} of positive integers chosen proportionally to yield the same power of L in the respective isomorphisms

(A.1) with $H^0(X', \mathcal{O}_{X'}(n))$ and $H^0(\tilde{X}', \mathcal{O}_{\tilde{X}'}(\tilde{n}))$. Taking, as powers of L , two sufficiently large consecutive multiples of the exponent of $M(X, L)$ yields respective line bundles $\mathcal{O}_{X'}(1)$, such that the pull-back of their difference is isomorphic to L^d , where d denotes the exponent of $M(X, L)$. If a line bundle L' on X' satisfies $\pi^*L' \cong L^e$ for some $e > 0$ then L' is ample, so all sufficiently large powers of L' are globally generated and thus e is divisible by d . \square

A morphism satisfying the properties stated in (i) is called a *contraction* and is necessarily surjective with geometrically connected fibers [31, 4.3.2, 4.3.4]. We call this the *contraction determined by L* .

Below we present a result which encompasses the following examples.

Example A.2. We suppose X is smooth and projective over k of dimension n , with ample line bundle L_0 , and effective divisor $E \subset X$.

- (i) If $E \cong \mathbb{P}^{n-1}$ with normal bundle $N_{E/X} \cong \mathcal{O}_{\mathbb{P}^{n-1}}(-1)$, and m denotes the positive integer with $L_0|_E \cong \mathcal{O}_{\mathbb{P}^{n-1}}(m)$, then $L := L_0(mE)$ is semiample and determines a contraction

$$X \cong B\ell_{\{p\}}X' \rightarrow X'$$

of E to a k -point p on a nonsingular projective variety X' .

- (ii) If $n = 3$ and $E \cong \mathbb{P}^1 \times \mathbb{P}^1$, with $N_{E/X} \cong \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-1, -1)$ and $L_0|_E \cong \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(m_1, m_2)$, then $L := L_0(\min(m_1, m_2)E)$ is semiample and determines a contraction

$$X \cong B\ell_C X' \rightarrow X'$$

for a curve $C \cong \mathbb{P}^1$ on nonsingular X' with $N_{C/X'} \cong \mathcal{O}_{\mathbb{P}^1}(-1)^2$, when $m_1 \neq m_2$, respectively

$$X \cong B\ell_{\{p\}}X' \rightarrow X'$$

for a k -point $p \in X'$ with an ordinary double point singularity, when $m_1 = m_2$.

When $\dim(X) = 2$, Example A.2(i) is the contraction theorem of Castelnuovo and Enriques and leads to minimal models of algebraic surfaces. The case of dimension at least 3 requires the sophistication of the minimal model program (see [41] for a survey), including the observation that a curve class (modulo numerical equivalence), which meets a canonical divisor positively and is extremal in the cone of effective classes, uniquely determines a contraction. Mori, in dimension 3, identifies five kinds of divisorial contractions [60, Thm. 3.3], including the ones given in Example A.2; we see already the possibility of contraction to a singular point, which is essential for the theory. The results below on divisorial contractions to a point (Proposition A.3) and to a general subscheme

(Proposition A.7) are essentially contained in [60, Lem. 3.32]; a relative version is presented in Proposition A.9.

An important ingredient, already appearing in loc. cit., is the following notion [61]: an ample line bundle L on a projective scheme X is *normally generated* if

$$\mathrm{Sym}^n H^0(X, L) \rightarrow H^0(X, L^n)$$

is surjective for every positive integer n .

Proposition A.3. *Let X be a projective scheme over k with ample line bundle L_0 and effective Cartier divisor $E \subset X$ with $H^0(E, \mathcal{O}_E) = k$ (this is the case, e.g., when geometrically E is connected and reduced). Suppose that $N_{E/X}^\vee$ is ample, with $L_0|_E \otimes (N_{E/X})^m$ a torsion line bundle for some $m > 0$. Assume, furthermore, that $N_{E/X}^\vee$ is normally generated, and $H^1(E, (N_{E/X}^\vee)^n) = 0$ for all $n > 0$. Then $L := L_0(mE)$ is semiample with exponent of $M(X, L)$ equal to the order of $L_0|_E \otimes (N_{E/X})^m$ in $\mathrm{Pic}(E)$ and determines a contraction $X \cong \mathrm{Bl}_{\{p\}}Y \rightarrow Y$ of E to a k -point $p \in Y$ such that the local ring $\mathcal{O}_{Y,p}$ has associated graded k -algebra isomorphic to $\bigoplus H^0(E, (N_{E/X}^\vee)^n)$.*

Proof. Replacing L_0 by some power, we may suppose L_0 is very ample with $H^1(X, L_0) = 0$, the restriction of L to E is trivial, and $m \geq 2$. Now, we claim, $H^1(X, L_0(jE)) = 0$ for $0 \leq j < m$. We have this for $j = 0$, and we argue by induction on j using the exact sequence in cohomology associated with

$$0 \rightarrow L_0((j-1)E) \rightarrow L_0(jE) \rightarrow (N_{E/X}^\vee)^{m-j} \rightarrow 0.$$

Then, by considering the sequence for $j = m$ we find that L has a global section which restricts to a nonzero constant on E (under $L|_E \cong \mathcal{O}_E$). Furthermore, by the sequence for $j = m-1$ we may lift a basis of $H^0(E, N_{E/X}^\vee)$ to global sections of L , vanishing along E . We may view, as well, all the global sections of L_0 as global sections of L . So we see that L is globally generated, and the induced morphism to its image Y in projective space contracts E to a point p , such that the scheme-theoretic pre-image of p is E , and $X \setminus E \cong Y \setminus \{p\}$. Let us denote by \mathcal{I}_E the ideal sheaf of E , so $\mathcal{I}_E/\mathcal{I}_E^2 \cong N_{E/X}^\vee$. Then we see from these two cases of the sequence,

$$\mathfrak{m}_p/\mathfrak{m}_p^2 \rightarrow H^0(E, \mathcal{I}_E/\mathcal{I}_E^2)$$

is surjective. Again replacing L_0 with a tensor power we may suppose that the morphism $\pi: X \rightarrow Y$ given by the global sections of L induces an isomorphism $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$. Then by the Theorem on Formal Functions, we have an isomorphism

$$\widehat{\mathcal{O}}_{Y,p} \xrightarrow{\sim} \varprojlim H^0(X, \mathcal{O}_X/\mathcal{I}_E^n). \tag{A.2}$$

We claim, the isomorphism (A.2) comes from isomorphisms

$$\mathcal{O}_{Y,p}/\mathfrak{m}_p^n \rightarrow H^0(X, \mathcal{O}_X/\mathcal{I}_E^n)$$

for every n .

We begin by observing, using the H^1 -vanishing hypothesis, that

$$H^0(X, \mathcal{I}_E^i/\mathcal{I}_E^{j+1}) \rightarrow H^0(X, \mathcal{I}_E^i/\mathcal{I}_E^j) \quad (\text{A.3})$$

is surjective for all $i < j$. We will use this fact several times in the remainder of the proof.

By the normal generation hypothesis, for every j , the homomorphism

$$H^0(E, \mathcal{I}_E/\mathcal{I}_E^2) \otimes H^0(E, \mathcal{I}_E^{j-1}/\mathcal{I}_E^j) \rightarrow H^0(E, \mathcal{I}_E^j/\mathcal{I}_E^{j+1}),$$

given by multiplication, is surjective. Combining this with the observation about $\mathfrak{m}_p/\mathfrak{m}_p^2$, we have the surjectivity of

$$\mathfrak{m}_p/\mathfrak{m}_p^2 \otimes H^0(E, \mathcal{I}_E^{j-1}/\mathcal{I}_E^j) \rightarrow H^0(E, \mathcal{I}_E^j/\mathcal{I}_E^{j+1}).$$

We see more generally that if s_1, \dots, s_ℓ denote lifts of a basis of $\mathfrak{m}_p/\mathfrak{m}_p^2$ to the coordinate ring of an affine neighborhood of p , then multiplication by s_1, \dots, s_ℓ induces surjective

$$H^0(X, \mathcal{I}_E^i/\mathcal{I}_E^j)^\ell \rightarrow H^0(X, \mathcal{I}_E^{i+1}/\mathcal{I}_E^{j+1}) \quad (\text{A.4})$$

for all $i < j$. Indeed, we have the exact sequences

$$0 \rightarrow H^0(X, \mathcal{I}_E^{j-1}/\mathcal{I}_E^j)^\ell \rightarrow H^0(X, \mathcal{I}_E^i/\mathcal{I}_E^j)^\ell \rightarrow H^0(X, \mathcal{I}_E^i/\mathcal{I}_E^{j-1})^\ell \rightarrow 0 \quad (\text{A.5})$$

and

$$0 \rightarrow H^0(X, \mathcal{I}_E^j/\mathcal{I}_E^{j+1}) \rightarrow H^0(X, \mathcal{I}_E^{i+1}/\mathcal{I}_E^{j+1}) \rightarrow H^0(X, \mathcal{I}_E^{i+1}/\mathcal{I}_E^j) \rightarrow 0, \quad (\text{A.6})$$

with compatible vertical maps given by multiplication with s_1, \dots, s_ℓ . Induction on j for each fixed i establishes the surjectivity of (A.4).

If we regard i as fixed and apply to (A.4) the general fact that an inverse system of surjective linear maps of finite-dimensional vector spaces induces a surjective map of inverse limits, then we find,

$$\mathfrak{m}_p \varprojlim H^0(X, \mathcal{I}_E^i/\mathcal{I}_E^n) = \varprojlim H^0(X, \mathcal{I}_E^{i+1}/\mathcal{I}_E^n). \quad (\text{A.7})$$

An inductive argument now shows that we have an isomorphism

$$\mathfrak{m}_p^i \widehat{\mathcal{O}}_{Y,p} \xrightarrow{\sim} \varprojlim H^0(X, \mathcal{I}_E^i/\mathcal{I}_E^n), \quad (\text{A.8})$$

for every i . The case $i = 0$ is (A.2), and from (A.8) we have an isomorphism of the respective ideals generated on each side by \mathfrak{m}_p , which with (A.7) gives us the inductive step.

The claim is now immediate from (A.2) and (A.8). Notice that the claim implies that for every n we have an isomorphism

$$\mathfrak{m}_p^{n-1}/\mathfrak{m}_p^n \rightarrow H^0(E, \mathcal{I}_E^{n-1}/\mathcal{I}_E^n). \quad (\text{A.9})$$

The statement concerning the associated graded ring of $\mathcal{O}_{Y,p}$ in the proposition is immediate from (A.9). By the universal property of blowing up, we obtain a unique morphism $X \rightarrow B\ell_{\{p\}}Y$ factorizing π , whose restriction to E is given by

$$E \cong \text{Proj} \left(\bigoplus H^0(E, \mathcal{I}_E^n/\mathcal{I}_E^{n+1}) \right) \xrightarrow{\sim} \text{Proj} \left(\bigoplus \mathfrak{m}_p^n/\mathfrak{m}_p^{n+1} \right).$$

Now $X \rightarrow B\ell_{\{p\}}Y$ is a morphism of projective schemes, such that the scheme-theoretic pre-image of every point of $B\ell_{\{p\}}Y$ is a reduced point with the same residue field. By [32, 18.12.6], this must be a closed immersion. But the exceptional divisor of the blow-up is an effective Cartier divisor, over the complement of which we have an isomorphism, hence $X \rightarrow B\ell_{\{p\}}Y$ is an isomorphism. \square

For the version of Proposition A.3 which allows contraction to a more general variety than a point, we need relative versions of the hypotheses. A relative version of the hypothesis $H^0(E, \mathcal{O}_E) = k$ is to be given a flat contraction morphism $E \rightarrow F$, and the conclusion will be that this is determined by $L|_E$.

Lemma A.4. *Let $Y = \text{Spec}(A)$ be a Noetherian affine scheme, $\pi: X \rightarrow Y$ a flat projective morphism, L an ample line bundle on X , and $y \in Y$ a point such that the restriction L_y of L to the fiber X_y is very ample (resp. is normally generated) and satisfies $H^1(X_y, L_y^n) = 0$, for $n = 1$ (resp. for all $n > 0$). Then there exists an open subscheme $U \subset Y$ with $y \in U$ such that relative to $X \times_Y U \rightarrow U$ the restriction of L is very ample (resp. is very ample with*

$$\text{Sym}^n H^0(X \times_Y U, L) \rightarrow H^0(X \times_Y U, L^n)$$

surjective for all $n > 0$.

Proof. By the machinery of cohomology and base change, we may replace Y by a Zariski neighborhood of y , such that the coherent sheaf π_*L is locally free and its formation commutes with arbitrary base change. Since L_y is globally generated, the cokernel of $\pi^*\pi_*L \rightarrow L$ has trivial fiber over y , hence may be assumed to vanish if we shrink Y further. We obtain a morphism $X \rightarrow \mathbb{P}((\pi_*L)^\vee)$ over Y , and this restricts to a closed immersion of X_y since L_y is very ample. The support of the cokernel of $\mathcal{O}_{\mathbb{P}_Y^N} \rightarrow \mathcal{O}_X$ is disjoint from X_y , hence by further shrinking we find that L is relatively very ample. There exists n_0 such that

$$\text{Sym}^n H^0(X, L) \rightarrow H^0(X, L^n)$$

is surjective for all $n > n_0$ (consequence of the vanishing of H^1 of all sufficiently large twists of the ideal sheaf of X in $\mathbb{P}(H^0(X, L)^\vee)$). Now suppose that L_y is normally generated. Then, as above, we may suppose that $\pi_* L^n$ is locally free and its formation commutes with arbitrary base change for all $1 \leq n \leq n_0$. For all such n , we consider the cokernel of $\text{Sym}^n H^0(X, L) \rightarrow H^0(X, L^n)$, which may be assumed to vanish by shrinking Y , and thus we have the desired conclusion when L_y is normally generated. \square

Lemma A.5. *Let X be a scheme, with projective morphism $\pi: X \rightarrow S$ to a Noetherian scheme S and relatively ample line bundles L and M . Then for every coherent sheaf \mathcal{F} on X there exists $n \in \mathbb{N}$, such that $R^i\pi_*(\mathcal{F} \otimes L^a \otimes M^b) = 0$ for all $i > 0$ and $a, b \in \mathbb{N}$ with $\max(a, b) \geq n$.*

Proof. We immediately reduce to the case S when is affine. Using the ampleness of L and M , we reduce further to establishing the assertion with $\max(a, b)$ replaced by $\min(a, b)$. By considering \mathcal{F} twisted by various powers of L and M and replacing L and M with suitable powers, we may suppose that L and M are very ample. By pushing forward \mathcal{F} along the closed immersion $X \rightarrow \mathbb{P}_S^\ell \times_S \mathbb{P}_S^m$ for some $\ell, m \in \mathbb{N}$ determined by the sections of L and M , we are reduced to proving the following assertion: if \mathcal{F} is a coherent sheaf on $\mathbb{P}_S^\ell \times_S \mathbb{P}_S^m$ then there exists $n \in \mathbb{N}$ such that

$$H^i(\mathbb{P}_S^\ell \times_S \mathbb{P}_S^m, \mathcal{F}(a, b)) = 0 \quad \text{for all } a, b \in \mathbb{N} \text{ with } \min(a, b) \geq n.$$

Here, $\mathcal{F}(a, b)$ denotes the twist by the pullback of $\mathcal{O}_{\mathbb{P}_S^\ell}(a)$ and the pullback of $\mathcal{O}_{\mathbb{P}_S^m}(b)$. The proof is by descending induction on i , starting from the case $i = \ell + m + 1$, which is trivial by the vanishing of cohomology in degrees greater than the relative dimension. For the inductive step, the ampleness of $\mathcal{O}_{\mathbb{P}_S^\ell \times_S \mathbb{P}_S^m}(1, 1)$ implies the existence of $c, d \in \mathbb{N}$ such that there is an exact sequence

$$0 \rightarrow \mathcal{G} \rightarrow \mathcal{O}_{\mathbb{P}_S^\ell \times_S \mathbb{P}_S^m}(-c, -c)^{\oplus d} \rightarrow \mathcal{F} \rightarrow 0$$

of coherent sheaves. Twisting and applying the induction hypothesis and known cohomology of line bundles on $\mathbb{P}_S^\ell \times_S \mathbb{P}_S^m$, we obtain the result. \square

Lemma A.6. *Let X be a scheme, with projective morphism $\pi: X \rightarrow S$ to a Noetherian scheme S and line bundles L and M , such that locally on S the line bundle L is semiample, determining a flat contraction $\psi: X \rightarrow Y$, and M is relatively ample. Suppose $c, d \in \mathbb{N}$ are such that for all $y \in Y$ the fiber X_y satisfies $H^i(X_y, (M|_{X_y})^j) = 0$ for all $1 \leq i \leq c$ and $j \geq d$. Then there exists $n \in \mathbb{N}$ such that*

$$R^i\pi_*(L^a \otimes M^b) = 0$$

for all $1 \leq i \leq c$ and $a, b \in \mathbb{N}$ such that $b \geq d$ and $\max(a, b) \geq n$.

Proof. As in the proof of the previous lemma, we may assume that S is affine and observe that the pair of maps to projective spaces, determined by suitable powers of L and M , leads to the vanishing of all higher cohomology for $\min(a, b) \geq n$, for some $n \in \mathbb{N}$. By ampleness, after possibly increasing n , we obtain the vanishing for all $a, b \in \mathbb{N}$ with $b \geq n$. Now we consider a fixed b with $d \leq b < n$ and e with $0 \leq e < \ell$, where ℓ is the exponent of $M(X, L)$, and apply the hypothesis, the Leray spectral sequence, and the machinery of cohomology and base change to obtain

$$H^i(X, L^{a\ell+e} \otimes M^b) \cong H^i(Y, \mathcal{O}_Y(a) \otimes \psi_*(L^e \otimes M^b)),$$

which by ampleness vanishes for sufficiently large a . This gives the result. \square

We recall [32, 6.10.1], a locally Noetherian scheme Y is said to be *normally flat* along a closed subscheme F if $\mathcal{I}_F^j/\mathcal{I}_F^{j+1}$ is a locally free sheaf of \mathcal{O}_F -modules for every j , where \mathcal{I}_F denotes the ideal sheaf of F .

Proposition A.7. *Let X be a projective scheme over k with ample line bundle L_0 and effective Cartier divisor $E \subset X$. Suppose for some $m > 0$, the line bundle $L_0|_E \otimes (N_{E/X})^m$ is semiample and determines a flat contraction $\psi: E \rightarrow F$. Suppose, furthermore, $N_{E/X}^\vee$ is ample relative to ψ and on fibers is normally generated with vanishing H^1 of all positive powers. Then $L := L_0(mE)$ is semiample with exponent of $M(X, L)$ equal to exponent of $M(E, L_0|_E \otimes (N_{E/X})^m)$ and determines a contraction $X \cong B\ell_F Y \rightarrow Y$ of E to F such that Y is normally flat along F with normal cone*

$$\text{Spec} \left(\bigoplus_{n \geq 0} \psi_*(N_{E/X}^\vee)^n \right).$$

Proof. Replacing L_0 by some power, we may suppose that L_0 is very ample, with $H^1(X, L_0) = 0$, $m \geq 2$, and $L|_E$ globally generated, with sections defining $\psi: E \rightarrow F$ as in the statement. Since the tensor product of a very ample line bundle and a globally generated line bundle is very ample [30, 4.4.8], for every $1 \leq a \leq m-1$ the line bundle

$$(L_0|_E)^{m-a} \otimes (L_0|_E \otimes N_{E/X}^m)^a \cong (L_0|_E)^m \otimes (N_{E/X})^{am}$$

is very ample. So, $L_0|_E \otimes (N_{E/X})^a$ is ample for $1 \leq a \leq m-1$. Now we claim,

$$H^1(E, (L_0|_E)^{n_1} \otimes (N_{E/X})^{n_2}) = 0$$

for all sufficiently large n_1 and $0 \leq n_2 < mn_1$. Indeed, Lemma A.5 applied to the pairs of line bundles $L_0|_E \otimes (N_{E/X})^{a-1}$ and $L_0|_E \otimes (N_{E/X})^a$ for $a = 1, \dots, m-1$ takes care of the case $0 \leq n_2 \leq (m-1)n_1$. We

consider the pair of line bundles $L|_E$ (semiample) and $M := L_0|_E \otimes (N_{E/X})^{m-1}$ (ample). Since the restriction of L to any fiber of ψ is trivial, the restriction of M is isomorphic to the restriction of $N_{E/X}^\vee$, and by assumption H^1 of all positive powers vanish. An application of Lemma A.6 completes the verification of the claim. As a consequence, after again replacing L_0 by a power, we may suppose that

$$H^1(E, L_0|_E \otimes (N_{E/X})^j) = 0 \quad (\text{A.10})$$

for $0 \leq j < m$.

The exact sequence

$$0 \rightarrow L_0((j-1)E) \rightarrow L_0(jE) \rightarrow L_0|_E \otimes (N_{E/X})^j \rightarrow 0$$

and an inductive argument lead to $H^1(X, L_0(jE)) = 0$ for $0 \leq j < m$. By the vanishing for $j = m-1$ and $j = m-2$, the homomorphisms

$$H^0(X, L) \rightarrow H^0(E, L|_E) \quad \text{and} \quad H^0(X, L(-E)) \rightarrow H^0(E, L|_E \otimes N_{E/X}^\vee)$$

are surjective. It follows that L is globally generated, and if we let $\pi: X \rightarrow Y$ denote the contraction that it determines, then the image of the restriction of π to E is a copy of F in Y , such that the scheme-theoretic pre-image of F under π is E . Moreover, π induces a surjective homomorphism of sheaves

$$\mathcal{I}_F/\mathcal{I}_F^2 \rightarrow \psi_*(\mathcal{I}_E/\mathcal{I}_E^2). \quad (\text{A.11})$$

Let \mathfrak{Y} denotes the formal completion of Y along F . By the Theorem on Formal Functions in the form [31, 4.1.5], π induces an isomorphism

$$\mathcal{O}_{\mathfrak{Y}} \xrightarrow{\sim} \varprojlim \pi_*(\mathcal{O}_X/\mathcal{I}_E^n). \quad (\text{A.12})$$

We claim that the isomorphism (A.12) comes from isomorphisms

$$\mathcal{O}_Y/\mathcal{I}_F^n \xrightarrow{\sim} \pi_*(\mathcal{O}_X/\mathcal{I}_E^n), \quad (\text{A.13})$$

for every n .

We start by observing, by the machinery of cohomology and base change, that the hypothesis of vanishing of H^1 of positive powers of $N_{E/X}^\vee$ implies $R^1\psi_*(\mathcal{I}_E^j/\mathcal{I}_E^{j+1}) = 0$ for all $j > 0$. A consequence is the surjectivity of

$$\pi_*(\mathcal{I}_E^i/\mathcal{I}_E^{i+1}) \rightarrow \pi_*(\mathcal{I}_E^i/\mathcal{I}_E^j) \quad (\text{A.14})$$

for all $i < j$. As a consequence, in the inverse system in (A.12) the transition maps are epimorphisms, and the same holds with \mathcal{O}_X replaced by any positive power of \mathcal{I}_E .

By Lemma A.4, the homomorphism

$$\psi_*(\mathcal{I}_E/\mathcal{I}_E^2) \otimes \psi_*(\mathcal{I}_E^{j-1}/\mathcal{I}_E^j) \rightarrow \psi_*(\mathcal{I}_E^j/\mathcal{I}_E^{j+1})$$

induced by π is surjective for all j . Combining this with (A.11) we have the surjectivity of

$$\mathcal{I}_F/\mathcal{I}_F^2 \otimes \psi_*(\mathcal{I}_E^{j-1}/\mathcal{I}_E^j) \rightarrow \psi_*(\mathcal{I}_E^j/\mathcal{I}_E^{j+1}).$$

The remainder of the argument follows closely the proof of Proposition A.3. If $V \subset Y$ is any affine open and s_1, \dots, s_ℓ are sections that generate $\mathcal{I}_F|_V$ as a sheaf of \mathcal{O}_V -modules, then multiplication by s_1, \dots, s_ℓ induces surjective

$$H^0(U, \mathcal{I}_E^i \cdot \mathcal{I}_E^j)^\ell \rightarrow H^0(U, \mathcal{I}_E^{i+1}/\mathcal{I}_E^{j+1}) \quad (\text{A.15})$$

for all $i < j$, where $U \subset X$ denotes the pre-image of V . We see this by considering exact sequences, analogous to (A.5)–(A.6), but with sections on U instead of X .

Fixing i and letting K_j denote the kernel of (A.15), we write the corresponding short exact sequences for consecutive values $j = n - 1$ and $j = n$ and connect them by homomorphisms of the form (A.14), to see that the induced homomorphism of kernels $K_n \rightarrow K_{n-1}$ is surjective. After applying \varprojlim we obtain a short exact sequence, which tells us that

$$\mathcal{I}_F \varprojlim \pi_*(\mathcal{I}_E^i/\mathcal{I}_E^n) = \varprojlim \pi_*(\mathcal{I}_E^{i+1}/\mathcal{I}_E^n).$$

An inductive argument, as in the proof of Proposition A.3, shows that we have an isomorphism

$$\mathcal{I}_F^i \mathcal{O}_{\mathfrak{Y}} \xrightarrow{\sim} \varprojlim \pi_*(\mathcal{I}_E^i/\mathcal{I}_E^n), \quad (\text{A.16})$$

for every i .

The claim is now immediate from (A.12) and (A.16). The claim implies that for every n we have an isomorphism

$$\mathcal{I}_F^{n-1}/\mathcal{I}_F^n \xrightarrow{\sim} \psi_*(\mathcal{I}_E^{n-1}/\mathcal{I}_E^n). \quad (\text{A.17})$$

The sheaf on the right is locally free by cohomology and base change, hence Y is normally flat along F . The statement concerning the normal cone in the proposition is immediate from (A.17). The conclusion of the proof is exactly as in Proposition A.3. \square

The general result and examples are valid as well in a relative setting. If S is a Noetherian scheme and X is a scheme, proper over S , then we call a line bundle L on X *relatively semiample* if over every affine open subscheme of S some positive power of L is globally generated. Equivalently, there exists $m > 0$ such that over every affine open subscheme L^m is globally generated. The set of natural numbers m with this property will be denoted by $M(X/S, L)$, or just $M(X, L)$ when no confusion will arise. There is, as before, a well-defined exponent of $M(X/S, L)$.

Proposition A.8. *Let X be a scheme with a proper morphism $\varphi: X \rightarrow S$ to a Noetherian scheme S , relatively semiample line bundle L , and surjective morphism of sheaves $\varphi^*\mathcal{E} \rightarrow L^m$, for some $m > 0$, and coherent sheaf \mathcal{E} on S , thereby inducing a morphism*

$$\psi: X \rightarrow \text{Proj}(\text{Sym}^\bullet \mathcal{E}).$$

Let $Y = \psi(X)$, and write the Stein factorization of ψ as proper morphism

$$\pi: X \rightarrow X',$$

followed by finite morphism $f: X' = \text{Spec}(\psi_\mathcal{O}_X) \rightarrow Y$. We let $\mathcal{O}_Y(1)$, respectively $\mathcal{O}_{X'}(1)$ denote the restriction to Y , respectively the pull-back to X' of $\mathcal{O}_{\text{Proj}(\text{Sym}^\bullet \mathcal{E})}(1)$, so $\pi^*\mathcal{O}_{X'}(1) \cong L^m$. Then:*

- (i) *The morphism $\pi: X \rightarrow X'$ is proper and induces an isomorphism $\mathcal{O}_{X'} \xrightarrow{\sim} \pi_*\mathcal{O}_X$.*
- (ii) *The line bundle $\mathcal{O}_{X'}(n)$ is relatively ample over S , and for any positive integer n such that $\mathcal{O}_{X'}(n)$ is relatively very ample, the morphism $X \rightarrow \text{Proj}(\text{Sym}^\bullet \varphi_*(L^{mn}))$ determined by the direct image of L^{mn} factors uniquely through X' , identifying X' with the image of X in $\text{Proj}(\text{Sym}^\bullet \varphi_*(L^{\otimes n}))$.*
- (iii) *With the property in (ii) the scheme X' , with $\pi: X \rightarrow X'$, is determined uniquely up to canonical isomorphism, independently of the choice of m , coherent sheaf \mathcal{E} , and morphism $\varphi^*\mathcal{E} \rightarrow L^m$, and the powers of L that are isomorphic to the pull-back of a line bundle from X' are precisely all multiples of the exponent of $M(X/S, L)$.*
- (iv) *If φ is flat and there is a positive integer n such that for all $i > 0$,*

$$R^i\varphi_*(L^n) = R^i\varphi_*(L^{2n}) = \dots = 0,$$

then the corresponding direct image sheaves are locally free, their formation commutes with arbitrary base change $S' \rightarrow S$, when S' is Noetherian the pull-back of L to $S' \times_S X$ determines the contraction $S' \times_S X \rightarrow S' \times_S X'$, and X' is flat over S .

Proof. As in Proposition A.1, (i) recalls a basic property of the Stein factorization. For (ii), we have as before the first statement by standard facts on ample line bundles. Letting $\rho: X' \rightarrow S$ denote the structure morphism, using the isomorphism $\mathcal{O}_{X'} \cong \pi_*\mathcal{O}_X$ we have $\varphi_*(L^{\otimes n}) \cong \rho_*(\mathcal{O}_{X'}(n))$, and this determines compatible morphisms to its projectivization from X and from X' , the latter a closed immersion. For (iii) the argument is as in Proposition A.1. The statements concerning the direct image sheaves in (iv) follow from cohomology and base change, and the flatness from

$$X' \cong \text{Proj}(\mathcal{O}_S \oplus \varphi_*(L^{\otimes n}) \oplus \varphi_*(L^{\otimes 2n}) \oplus \dots).$$

The remaining assertion follows from these facts, using that some multiple of n satisfies the condition in (ii) both for X and for $S' \times_S X$. \square

Proposition A.9. *Let S be a Noetherian scheme, and let X be a projective scheme over S (resp. a projective flat scheme over S) with relatively ample line bundle L_0 and effective Cartier divisor $E \subset X$. Suppose for some $m > 0$, the line bundle $L_0|_E \otimes (N_{E/X})^m$ over affine open subsets of S is semiample and determines a flat contraction $\psi: E \rightarrow F$. Suppose, furthermore, $N_{E/X}^\vee$ is ample relative to ψ and on fibers is normally generated with vanishing H^1 of all positive powers (resp. vanishing H^i for all $i > 0$ of all nonnegative powers). Then $L := L_0(mE)$ is semiample with exponent of $M(X, L)$ equal to exponent of $M(E, L_0|_E \otimes (N_{E/X})^m)$ over affine open subsets of S and determines a contraction $X \cong Bl_F Y \rightarrow Y$ of E to F (resp. a contraction $X \cong Bl_F Y \rightarrow Y$ of E to F with Y flat over S , whose formation commutes with arbitrary base change $S' \rightarrow S$ with S' Noetherian) such that Y is normally flat along F with normal cone*

$$\text{Spec} \left(\bigoplus_{n \geq 0} \psi_*(N_{E/X}^\vee)^n \right)$$

Proof. We may suppose S is affine. Now we argue as in the proof of Proposition A.7, relative over S . In the flat case with the stronger hypothesis of cohomology vanishing we have (A.10) also for $j = m$, also with L_0 replaced by L_0^n for any $n > 0$ and all $0 \leq j \leq mn$, and also with any H^i ($i > 0$) in place of H^1 . Hence $H^i(X, L^n) = 0$ for all $i > 0$ and $n > 0$. Proposition A.8(iv) gives us the desired conclusion. \square

Remark A.10. Proposition A.9 is a strict generalization of Proposition A.7, which in turn generalizes Proposition A.3; the inclusion of all three results serves expository purposes only.

Remark A.11. In Proposition A.3, respectively A.7 and A.9, the morphism $\pi: X \rightarrow Y$ induces by pullback and direct image an equivalence of categories between locally free coherent sheaves on Y and locally free coherent sheaves on X whose restriction to E is free, respectively isomorphic to ψ^* of a locally free coherent sheaf on F . That π_* of such a sheaf \mathcal{F} is locally free follows from the Theorem on Formal Functions and observations concerning $H^0(X, \mathcal{I}_E^i \mathcal{F} / \mathcal{I}_E^j \mathcal{F})$, respectively $H^0(U, \mathcal{I}_E^i \mathcal{F} / \mathcal{I}_E^j \mathcal{F})$, analogous to those in (A.3)–(A.4), respectively (A.14)–(A.15).

Remark A.12. In Proposition A.9 (without the assumption that $X \rightarrow S$ is flat and stronger cohomology vanishing from the statement), $\pi: X \rightarrow Y$ is characterized by the following universal property: for any S -scheme Z , S -morphisms $Y \rightarrow Z$ are by composition with π in bijection with

S -morphisms $X \rightarrow Z$ whose restriction to E factors through F (by ψ); cf. [30, 8.11.1]. Here we may allow Z to be an algebraic space, in which case the verification uses the observation that given compatible separated étale morphisms of schemes $X' \rightarrow X$, $E' \rightarrow E$, and $F' \rightarrow F$, the functor on Y -schemes that sends $\tilde{Y} \rightarrow Y$, determining $\tilde{X} \rightarrow X$, $\tilde{E} \rightarrow E$, and $\tilde{F} \rightarrow F$ by base change, to the set of compatible lifts $\tilde{X} \rightarrow X'$, $\tilde{E} \rightarrow E'$, and $\tilde{F} \rightarrow F'$, is representable by a scheme Y' , separated and étale over Y . Already the functor sending $\tilde{Y} \rightarrow Y$ to the set of lifts $\tilde{F} \rightarrow F'$ is representable by an algebraic space, étale over Y [62, §1.1] [6, §5.1], so we may suppose $F' = F$ (and $E' = E$). Then the observation follows from a general result of Artin on contractions [7, Thm. 3.1] and the property of separated étale morphisms of algebraic spaces that if the target is a scheme then so is the source [43, Cor. II.6.17]. The universal property in this level of generality compares nicely with earlier results, e.g., in [58].

REFERENCES

- [1] D. Abramovich, T. Graber, and A. Vistoli. Gromov-Witten theory of Deligne-Mumford stacks. *Amer. J. Math.*, 130(5):1337–1398, 2008.
- [2] D. Abramovich, M. Olsson, and A. Vistoli. Tame stacks in positive characteristic. *Ann. Inst. Fourier (Grenoble)*, 58(4):1057–1091, 2008.
- [3] N. Addington, B. Hassett, Yu. Tschinkel, and A. Várilly-Alvarado. Cubic fourfolds fibered in sextic del Pezzo surfaces, 2016. [arXiv:1606.05321](https://arxiv.org/abs/1606.05321).
- [4] J. Alper. Good moduli spaces for Artin stacks. *Ann. Inst. Fourier (Grenoble)*, 63(6):2349–2402, 2013.
- [5] J. Alper, J. Hall, and D. Rydh. A Luna étale slice theorem for algebraic stacks, 2015. [arXiv:1504.06467](https://arxiv.org/abs/1504.06467).
- [6] J. Alper and A. Kresch. Equivariant versal deformations of semistable curves. *Michigan Math. J.*, 65(2):227–250, 2016.
- [7] M. Artin. Algebraization of formal moduli. II. Existence of modifications. *Ann. of Math. (2)*, 91:88–135, 1970.
- [8] M. Artin. Brauer-Severi varieties. In *Brauer groups in ring theory and algebraic geometry (Wilrijk, 1981)*, volume 917 of *Lecture Notes in Math.*, pages 194–210. Springer, Berlin-New York, 1982.
- [9] M. Artin. Left ideals in maximal orders. In *Brauer groups in ring theory and algebraic geometry (Wilrijk, 1981)*, volume 917 of *Lecture Notes in Math.*, pages 182–193. Springer, Berlin-New York, 1982.
- [10] M. Artin. Local structure of maximal orders on surfaces. In *Brauer groups in ring theory and algebraic geometry (Wilrijk, 1981)*, volume 917 of *Lecture Notes in Math.*, pages 146–181. Springer, Berlin-New York, 1982.
- [11] M. Artin, A. Grothendieck, and J. L. Verdier. *Théorie des topos et cohomologie étale des schémas (SGA 4)*, tome 3, volume 305 of *Lecture Notes in Math.* Springer, Berlin-New York, 1973.
- [12] M. Artin and D. Mumford. Some elementary examples of unirational varieties which are not rational. *Proc. London Math. Soc. (3)*, 25:75–95, 1972.

- [13] A. Auel, M. Bernardara, and M. Bolognesi. Fibrations in complete intersections of quadrics, Clifford algebras, derived categories, and rationality problems. *J. Math. Pures Appl. (9)*, 102(1):249–291, 2014.
- [14] A. Auel, Chr. Böhning, and H.-Chr. Graf von Bothmer. The transcendental lattice of the sextic Fermat surface. *Math. Res. Lett.*, 20(6):1017–1031, 2013.
- [15] A. Beauville. A very general sextic double solid is not stably rational. *Bull. Lond. Math. Soc.*, 48(2):321–324, 2016.
- [16] D. Bergh. Functorial destackification of tame stacks with abelian stabilisers. *Compos. Math.*, 153(6):1257–1315, 2017.
- [17] S. Bloch and A. Ogus. Gersten’s conjecture and the homology of schemes. *Ann. Sci. École Norm. Sup. (4)*, 7:181–201 (1975), 1974.
- [18] C. Böhning and H.-Chr. Graf von Bothmer. On stable rationality of some conic bundles and moduli spaces of Prym curves, 2016. [arXiv:1605.03029](https://arxiv.org/abs/1605.03029), to appear in Comm. Math. Helvetici.
- [19] J. W. Bruce and C. T. C. Wall. On the classification of cubic surfaces. *J. London Math. Soc. (2)*, 19(2):245–256, 1979.
- [20] C. Cadman. Using stacks to impose tangency conditions on curves. *Amer. J. Math.*, 129(2):405–427, 2007.
- [21] J.-L. Colliot-Thélène. Birational invariants, purity and the Gersten conjecture. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 1–64. Amer. Math. Soc., Providence, RI, 1995.
- [22] J.-L. Colliot-Thélène and M. Ojanguren. Variétés unirationnelles non rationnelles: au-delà de l’exemple d’Artin et Mumford. *Invent. Math.*, 97(1):141–158, 1989.
- [23] J.-L. Colliot-Thélène and A. Pirutka. Hypersurfaces quartiques de dimension 3: non-rationalité stable. *Ann. Sci. École Norm. Sup. (4)*, 49(2):371–397, 2016.
- [24] A. J. de Jong. The period-index problem for the Brauer group of an algebraic surface. *Duke Math. J.*, 123(1):71–94, 2004.
- [25] P. Deligne and D. Mumford. The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.*, 36:75–109, 1969.
- [26] S. P. Diaz. Porteous’s formula for maps between coherent sheaves. *Michigan Math. J.*, 52(3):507–514, 2004.
- [27] D. Edidin, B. Hassett, A. Kresch, and A. Vistoli. Brauer groups and quotient stacks. *Amer. J. Math.*, 123(4):761–777, 2001.
- [28] W. Fulton. *Algebraic curves. An introduction to algebraic geometry*. W. A. Benjamin, Inc., New York-Amsterdam, 1969. Notes written with the collaboration of Richard Weiss, Mathematics Lecture Notes Series.
- [29] B. H. Gross and J. Harris. On some geometric constructions related to theta characteristics. In *Contributions to automorphic forms, geometry, and number theory*, pages 279–311. Johns Hopkins Univ. Press, Baltimore, MD, 2004.
- [30] A. Grothendieck. Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes. *Inst. Hautes Études Sci. Publ. Math.*, 8, 1961.
- [31] A. Grothendieck. Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. *Inst. Hautes Études Sci. Publ. Math.*, 11, 17, 1961–63.
- [32] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. *Inst. Hautes Études Sci. Publ. Math.*, 20, 24, 28, 32, 1964–67.

- [33] A. Grothendieck. Le groupe de Brauer. I–III. In *Dix exposés sur la cohomologie des schémas*, volume 3 of *Adv. Stud. Pure Math.*, pages 46–188. North-Holland, Amsterdam, 1968.
- [34] A. Grothendieck. Techniques de construction et théorèmes d’existence en géométrie algébrique. IV. Les schémas de Hilbert. In *Séminaire Bourbaki, Vol. 6*, pages Exp. No. 221, 249–276. Soc. Math. France, Paris, 1995.
- [35] B. Hassett, A. Kresch, and Yu. Tschinkel. On the moduli of degree 4 Del Pezzo surfaces. In *Development of moduli theory (Kyoto 2013)*, volume 69 of *Adv. Stud. Pure Math.*, pages 349–386. Math. Soc. Japan, Tokyo, 2016.
- [36] B. Hassett, A. Kresch, and Yu. Tschinkel. Stable rationality and conic bundles. *Math. Ann.*, 365(3-4):1201–1217, 2016.
- [37] B. Hassett, A. Kresch, and Yu. Tschinkel. Stable rationality in smooth families of threefolds, 2018. [arXiv:1802.06107](https://arxiv.org/abs/1802.06107).
- [38] B. Hassett, A. Pirutka, and Yu. Tschinkel. Stable rationality of quadric surface bundles over surfaces. *Acta Mathematica*, 220:341–365, 2018.
- [39] B. Hassett and Yu. Tschinkel. On stable rationality of Fano threefolds and del Pezzo fibrations. *J. Reine Angew. Math.*, 751:275–287, 2019.
- [40] H. Hironaka. Smoothing of algebraic cycles of small dimensions. *Amer. J. Math.*, 90:1–54, 1968.
- [41] Y. Kawamata, K. Matsuda, and K. Matsuki. Introduction to the minimal model problem. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 283–360. North-Holland, Amsterdam, 1987.
- [42] S. Keel and S. Mori. Quotients by groupoids. *Ann. of Math. (2)*, 145(1):193–213, 1997.
- [43] D. Knutson. *Algebraic spaces*. Lecture Notes in Mathematics, Vol. 203. Springer-Verlag, Berlin-New York, 1971.
- [44] M. Kontsevich and Yu. Tschinkel. Specialization of birational types, 2017. [arXiv:1708.05699](https://arxiv.org/abs/1708.05699).
- [45] A. Kresch. Hodge-theoretic obstruction to the existence of quaternion algebras. *Bull. London Math. Soc.*, 35(1):109–116, 2003.
- [46] A. Kresch. Flattening stratification and the stack of partial stabilizations of prestable curves. *Bull. Lond. Math. Soc.*, 45(1):93–102, 2013.
- [47] A. Kresch and Yu. Tschinkel. Stable rationality of Brauer-Severi surface bundles, 2017. [arXiv:1709.10151](https://arxiv.org/abs/1709.10151), to appear in *Manuscripta Math.*
- [48] A. Kresch and Yu. Tschinkel. Involution surface bundles over surfaces, 2018. [arXiv:1807.01592](https://arxiv.org/abs/1807.01592).
- [49] I. Krylov and T. Okada. Stable rationality of del Pezzo fibrations of low degree over projective spaces, 2017. [arXiv:1701.08372](https://arxiv.org/abs/1701.08372).
- [50] A. Kuznetsov. Derived categories of families of sextic del Pezzo surfaces, 2017. [arXiv:1708.00522](https://arxiv.org/abs/1708.00522).
- [51] G. Laumon and L. Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer-Verlag, Berlin, 2000.
- [52] R. Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer-Verlag, Berlin, 2004.
- [53] M. Lieblich. Twisted sheaves and the period-index problem. *Compos. Math.*, 144(1):1–31, 2008.
- [54] M. Lieblich. Period and index in the Brauer group of an arithmetic surface. *J. Reine Angew. Math.*, 659:1–41, 2011. With an appendix by Daniel Krashen.

- [55] T. Maeda. On standard projective plane bundles. *J. Algebra*, 197(1):14–48, 1997.
- [56] K. Matsuki and M. Olsson. Kawamata-Viehweg vanishing as Kodaira vanishing for stacks. *Math. Res. Lett.*, 12(2-3):207–217, 2005.
- [57] H. Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid.
- [58] J. Mazur. Conditions for the existence of contractions in the category of algebraic spaces. *Trans. Amer. Math. Soc.*, 209:259–265, 1975.
- [59] J. S. Milne. *Étale cohomology*, volume 33 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1980.
- [60] S. Mori. Threefolds whose canonical bundles are not numerically effective. *Ann. of Math. (2)*, 116(1):133–176, 1982.
- [61] D. Mumford. Varieties defined by quadratic equations. In *Questions on Algebraic Varieties (C.I.M.E., III Ciclo, Varenna, 1969)*, pages 29–100. Edizioni Cremonese, Rome, 1970.
- [62] A. M. Mustaţă and A. Mustaţă. The structure of a local embedding and Chern classes of weighted blow-ups. *J. Eur. Math. Soc.*, 14(6):1739–1794, 2012.
- [63] J. Nicaise and E. Shinder. The motivic nearby fiber and degeneration of stable rationality, 2017. [arXiv:1708.02790](https://arxiv.org/abs/1708.02790).
- [64] J. Oesinghaus. Conic bundles and iterated root stacks. *Eur. J. Math.*, 5(2):518–527, 2019.
- [65] M. Olsson and J. Starr. Quot functors for Deligne-Mumford stacks. *Comm. Algebra*, 31(8):4069–4096, 2003. Special issue in honor of Steven L. Kleiman.
- [66] M. Raynaud and L. Gruson. Critères de platitude et de projectivité. Techniques de “platification” d’un module. *Invent. Math.*, 13:1–89, 1971.
- [67] D. Rydh. Étale dévissage, descent and pushouts of stacks. *J. Algebra*, 331:194–223, 2011.
- [68] D. Rydh. Existence and properties of geometric quotients. *J. Algebraic Geom.*, 22(4):629–669, 2013.
- [69] D. J. Saltman. *Lectures on division algebras*, volume 94 of *CBMS Regional Conference Series in Mathematics*. Published by American Mathematical Society, Providence, RI; on behalf of Conference Board of the Mathematical Sciences, Washington, DC, 1999.
- [70] V. G. Sarkisov. On conic bundle structures. *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(2):371–408, 432, 1982.
- [71] V. V. Shokurov. Prym varieties: theory and applications. *Izv. Akad. Nauk SSSR Ser. Mat.*, 47(4):785–855, 1983.
- [72] M. G. Smith and J. S. Wilson. On subgroups of finite index in compact Hausdorff groups. *Arch. Math. (Basel)*, 80(2):123–129, 2003.
- [73] K.-O. Stöhr and J. F. Voloch. Weierstrass points and curves over finite fields. *Proc. London Math. Soc. (3)*, 52(1):1–19, 1986.
- [74] B. Totaro. Hypersurfaces that are not stably rational. *J. Amer. Math. Soc.*, 29(3):883–891, 2016.
- [75] C. Voisin. Unirational threefolds with no universal codimension 2 cycle. *Invent. Math.*, 201(1):207–237, 2015.
- [76] J. Włodarczyk. Simple Hironaka resolution in characteristic zero. *J. Amer. Math. Soc.*, 18(4):779–822, 2005.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE
190, CH-8057 ZÜRICH, SWITZERLAND

Email address: andrew.kresch@math.uzh.ch

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA

SIMONS FOUNDATION, 160 FIFTH AV., NEW YORK, NY 10010, USA

Email address: tschinkel@cims.nyu.edu