

# Conceptual dynamical models for turbulence

Andrew J. Majda\* and Yoonsang Lee\*

\*Department of Mathematics and Center for Atmosphere and Ocean Science, Courant Institute of Mathematical Sciences, New York University, New York, NY 10012

Contributed by Andrew J. Majda

Understanding the complexity of anisotropic turbulent processes in engineering and environmental fluid flows is a formidable challenge with practical significance since energy often flows intermittently from the smaller scales to impact the largest scales in these flows. Conceptual dynamical models for anisotropic turbulence are introduced and developed here which, despite their simplicity, capture key features of vastly more complicated turbulent systems. These conceptual models involve a large scale mean flow and turbulent fluctuations on a variety of spatial scales with energy conserving wave-mean flow interactions as well as stochastic forcing of the fluctuations. Numerical experiments with a six dimensional conceptual dynamical model confirm that these models capture key statistical features of vastly more complex anisotropic turbulent systems in a qualitative fashion. These features include chaotic statistical behavior of the mean flow with a sub-Gaussian probability distribution function (pdf) for its fluctuations while the turbulent fluctuations have decreasing energy and correlation times at smaller scales with nearly Gaussian pdfs for the large scale fluctuations and fat-tailed non-Gaussian pdfs for the smaller scale fluctuations. This last feature is a manifestation of intermittency of the small scale fluctuations where turbulent modes with small variance have relatively frequent extreme events which directly impact the mean flow. The dynamical modes introduced here potentially provide a useful test bed for algorithms for prediction, uncertainty quantification, and data assimilation for anisotropic turbulent systems.

intermittency | wave-mean interaction | stochastic model

Understanding the complexity of anisotropic turbulence processes over a wide range of spatiotemporal scales in engineering shear turbulence [1, 2, 3] as well as climate atmosphere ocean science [4, 5, 6] is a grand challenge of contemporary science. This is especially important from a practical viewpoint since energy often flows intermittently from the smaller scales to effect the largest scales in such anisotropic turbulent flows. The typical features of such anisotropic turbulent flows are the following [2, 3, 4]:

- (A) The large scale mean flow is usually chaotic but more predictable than the smaller scale fluctuations. The overall single point probability distribution function (pdf) of the flow field is nearly Gaussian while the mean flow pdf is sub-Gaussian, in other words, with less extreme variability than a Gaussian random variable.
- (B) There are nontrivial nonlinear interactions between the large scale mean flow and the smaller scale fluctuations which conserve energy.
- (C) There is a wide range of spatial scales for the fluctuations with features that the large scale components of the fluctuations contain more energy than the smaller scale components. Furthermore, these large scale fluctuating components decorrelate faster in time than the mean flow fluctuations on the largest scales while the smaller scale fluctuating components decorrelate faster in time than the larger scale fluctuating components.
- (D) The pdfs of the larger scale fluctuating components of the turbulent field are nearly Gaussian while the smaller scale fluctuating components are intermittent, and have fat tailed pdfs, in other words, a much higher probability of extreme events than a Gaussian distribution (see Figure 8.4 and 8.5 from [3] for such experimental features in a turbulent jet).

The goal here is to develop the simplest conceptual dynamical model for anisotropic turbulence that captures all of the features in (A)-(D) in a transparent qualitative fashion. In contrast to deterministic models of turbulence which are derived by Galerkin truncation of the Navier-Stokes equation [7] and do not display all the features in (A)-(D), the conceptual models developed here are low dimensional stochastic dynamical systems; the nonlinear interactions between the large scale mean flow component and the smaller scale fluctuating components are completely deterministic but the potential direct nonlinear interactions between the smaller scale fluctuating components are modeled stochastically by damping and stochastic forcing [6, 8]. The conceptual models developed here are not derived quantitatively from the Navier-Stokes equations but are developed to capture the key features in anisotropic turbulent flows listed in (A)-(D). by mimicking key physical processes. Besides aiding the understanding of anisotropic turbulent flows, such conceptual models are useful for designing and testing numerical algorithms for prediction and data assimilation in such complex turbulent systems.

## The Conceptual Model

The model has a mean scalar variable  $\bar{u}$ , representing the largest scales and a family of small scale variables,  $\vec{u}' = (u'_1, u'_2, \dots, u'_K)^T \in \mathbb{R}^K$  so that there are  $\mathbb{R}^{K+1}$  variables in the system  $\vec{u} = \begin{pmatrix} \bar{u} \\ \vec{u}' \end{pmatrix}$ . The variables  $u'_k, 1 \leq k \leq K$  represent contributions to the turbulent fluctuations from increasingly smaller scales as  $k$  increases with

$$u' = \sum_{k=1}^K u'_k, \quad [1]$$

the turbulent fluctuations. One can think of  $\bar{u}$  as the large scale spatial average of the turbulent dynamics at a single grid point in a more complex system and  $u'$  as the turbulent fluctuations at the grid point with

$$u(t) = \bar{u}(t) + \sum_{k=1}^K u'_k(t) \quad [2]$$

the total turbulent field. Note that the large scale mean  $\bar{u}$  can have fluctuating, chaotic dynamics in time through interactions with turbulence and its own intrinsic dynamics. The nonlinear interactions in turbulence conserve the total energy

Author contributions : A.J.M. designed research, Y.L. and A.J.M. performed research and analyzed data, A.J.M. wrote the paper

The authors declare no conflict of interest.

To whom correspondence may be addressed. E-mail : jonjon@cims.nyu.edu or ylee@cims.nyu.edu

of the mean and fluctuations and a key feature of the conceptual model is to utilize nonlinear interactions which conserve the energy  $E$ , which we take as given by

$$E(\bar{u}, \bar{u}') = \frac{1}{2}(\bar{u}^2 + \sum_{k=1}^K (u'_k)^2). \quad [3]$$

A hallmark of turbulence is that the large scales can destabilize the smaller scales in the turbulent fluctuations intermittently and this increased small scale energy can impact the large scales; this key feature is captured in the conceptual models. With the above discussion, here are the simplest models with all these features, the conceptual dynamical models for turbulence

$$\begin{aligned} \frac{d\bar{u}}{dt} &= -\bar{d}\bar{u} + \gamma \sum_{k=1}^K (u'_k)^2 - \bar{\alpha}\bar{u}^3 + \bar{F}, \\ \frac{du'_k}{dt} &= -d_k u'_k - \gamma \bar{u} u'_k + \sigma_k \dot{W}_k, \quad 1 \leq k \leq K. \end{aligned} \quad [4]$$

The reader can think of  $u'_k$  as the amplitude of the  $k$ -th Fourier cosine mode to aid the interpretation of the model but this is not necessary here. The system of  $K+1$  SDEs in [4] is written in physicist's notation with  $\dot{W}_k$  independent white noises for each  $k$  but the system in [4] is always interpreted in the Ito sense below. The reader easily verifies that the nonlinear interactions in [4] conserve the energy,  $E$ , in [3] which can be modified by the linear terms, the external forcing,  $\bar{F}$ , nonlinearity of the large scales, and the random forcing of the small scales. The turbulence dissipation coefficient  $d_k$  for  $k = 1, 2, \dots, K$  are positive,  $d_k > 0$ , in order for the turbulence to have a statistical steady state but the coefficient  $\bar{d}$ , for the large scales can be either positive or negative reflecting large scale instability; when  $\bar{d}$  is negative so there is instability on the large scales we add the stabilizing cubic term with  $\bar{\alpha} > 0$  while for positive  $\bar{d}$ , we assume  $\bar{\alpha} = 0$  and both cases are studied below. The external force,  $\bar{F}$  is a constant which is varied below to mimic fully turbulent regimes with (A)-(D). For a fixed coefficient of nonlinear interaction,  $\gamma > 0$ , there is local growth and instability in time for the  $k$ -th turbulent scale provided that

$$-d_k - \gamma \bar{u} > 0, \quad \text{i.e. } \bar{u} < \frac{-d_k}{\gamma} \quad [5]$$

and chaotic fluctuations of  $\bar{u}$  will create intermittency in  $u'_k$  through this mechanism. Thus, the overall system can have a statistical steady state while there is intermittent instability on the small scales which increases their energy and impacts the large scales creating non-Gaussian intermittent behavior in the system. With  $\bar{u} \equiv 0$ , the equation for the  $k$ -th turbulent scale,  $u'_k$ , is a simple Langevin process with Gaussian statistical steady state with zero mean and variance  $\frac{\sigma_k^2}{2d_k} = E_k$ ; it is natural to pick these energy densities to have power law behavior for this energy spectrum, i.e.,

$$\frac{\sigma_k^2}{2d_k} = E_k = E_0 |k|^{-\alpha}, \quad [6]$$

with  $E_0 > 0$  and  $\alpha \geq 0$  fixed constants [8]. For example,  $\alpha = \frac{5}{3}$  corresponds to the Kolmogorov spectrum [3, 8]. Note that we could allow coefficient  $\gamma$  in [4] to vary with  $k$  for  $k = 1, 2, \dots, K$  but we refrain from discussing this generalization here. On the other hand, it is natural to have the damping  $d_k$  vary with  $k$  to represent various dissipative processes such as viscosity or Ekman friction [8]. This completes the description of the conceptual models.

## Mathematical Properties

Note that the equation for the large scale mean,  $\bar{u}$ , is deterministic and without any direct stochastic forcing; this deterministic structure mimics that at the large scales for realistic turbulent flows. Nevertheless, the large scale mean  $\bar{u}$  interacts with the fluctuations  $u'_k$  which are stochastically forced. We claim that even with the above degenerate noise, the conceptual models in [4] are geometrically ergodic [9]; in other words, for any value of  $\bar{F}$ , a unique smooth ergodic invariant measure exists with exponential convergence of suitable statistics from time averages in the long time limit. To prove this, we apply the main theorem in [9] with the Lyapunov function given by the total energy in [3]. Two things need to be checked; the first is the coercivity of the generator applied to the Lyapunov function which is immediately satisfied given our hypotheses; the second condition is the hypoellipticity of the generator of [4]. To check hypoellipticity we consider the  $K$ -vector fields

$$X_k = (\sigma_k \delta_{ik}), \quad 0 \leq i \leq k, 1 \leq k \leq K, X_k \in \mathbb{R}^{K+1}, 1 \leq k \leq K$$

and

$$Y = \begin{pmatrix} -\bar{d}\bar{u} + \gamma \sum_{k=1}^K (u'_k)^2 - \bar{\alpha}\bar{u}^3 + \bar{F} \\ -d_k u'_k - \gamma \bar{u} u'_k \end{pmatrix}, \quad Y \in \mathbb{R}^{K+1}.$$

We only need to show that  $X_k, [X_k, Y], [X_k, [X_k, Y]]$  span all of  $\mathbb{R}^{K+1}$  where  $[X, Y] = X \cdot \nabla Y - Y \cdot \nabla X$  is the Lie bracket.

Since  $[X_k, [X_k, Y]] = \begin{pmatrix} 2\gamma\sigma_k^2 \\ 0 \end{pmatrix}$  and the  $X_k, 1 \leq k' \leq K$  span the orthogonal complement, hypoellipticity is satisfied.

## Phase Plane Analysis

Here we develop intuition regarding the parameters of the conceptual models which provide important guidelines to demonstrate below that these models with  $K \geq 2$  can capture all the features of anisotropic turbulence listed in (A)-(D) above. For such intuition, there is a revealing phase plane analysis of the two dimensional system for  $(\bar{u}, u')$  which is the special case of the model in [4] which  $K = 1$  and without noise. This system is given by

$$\begin{aligned} \frac{d\bar{u}}{dt} &= -\bar{d}\bar{u} + \gamma(u')^2 - \bar{\alpha}\bar{u}^3 + \bar{F}, \\ \frac{du'}{dt} &= -(d + \gamma\bar{u})u'. \end{aligned} \quad [7]$$

The linear subspace,  $(\bar{u}, 0)$ , is invariant for the dynamics which reduces on this subspace to the scalar equation

$$\frac{d\bar{u}}{dt} = -\bar{d}\bar{u} - \bar{\alpha}\bar{u}^3 + \bar{F} \quad [8]$$

while the general dynamics of [7] is invariant under the flip symmetry,  $(\bar{u}, u') \rightarrow (\bar{u}, -u')$ . Thus, there are between one and three critical points of [7] with the form  $(\bar{u}_{CR}, 0)$  as  $\bar{F}$  varies provided that  $\bar{d} < 0$  and  $\bar{\alpha} > 0$  and only a single critical point of the form  $(\bar{u}_{CR}, 0)$  with  $\bar{u}_{CR} = \frac{\bar{F}}{\bar{d}}$  for  $\bar{d} > 0, \bar{\alpha} = 0$ ; regardless of these stability properties along the  $\bar{u}$ -axis, such critical points are unstable to  $u'$  perturbations if and only if  $d + \gamma\bar{u}_{CR} < 0$ , i.e., the instability condition in [5] is satisfied. For suitable values of  $\bar{F}$ , there is another family of critical points for [7] with the form  $(\bar{u}_*, \pm u'_{CR})$  where

$$\begin{aligned} \bar{u}_* &= -\frac{d}{\gamma} \\ \gamma u_{CR}^2 &= \bar{d}\bar{u} + \bar{\alpha}\bar{u}^3 - \bar{F} \end{aligned} \quad [9]$$

Note that  $\bar{u}_*$  is exactly the critical value of neutral stability from [5] for the conceptual model. The linear stability matrix at these critical points for [7] has the form

$$\begin{pmatrix} -\bar{d} - 3\bar{\alpha}\bar{u}_*^2 & 2\omega \\ -\omega & 0 \end{pmatrix} \quad [10]$$

so these critical points are stable (unstable) if and only if  $-\bar{d} - 3\bar{\alpha}\bar{u}_*^2 < 0 (> 0)$ .

To develop guidelines in choosing parameters for the numerical experiments for  $K \geq 2$  with the conceptual model in [4], we consider the phase plane analysis in two scenarios with positive and negative large scale damping. In both cases, the parameter  $\gamma = 1.5$  and  $d = d_1 \equiv 1$  are fixed below while for

$$\begin{aligned} &\text{positive large scale damping, } \bar{d} = 0.01 \text{ and } \bar{\alpha} = 0 \\ &\text{negative large scale damping, } \bar{d} = -0.1 \text{ and } \bar{\alpha} = 0.05. \end{aligned} \quad [11]$$

First consider positive large scale damping; the two critical points  $(\bar{u}_*, \pm u'_{CR})$  occur for  $\bar{F} < \bar{F}_{CR} = \frac{-\bar{d}d}{\gamma} = -0.0067$  and are both stable by the criterion in [10] while the critical point  $(\frac{\bar{F}}{d}, 0)$  along the  $\bar{u}$ -axis is unstable to  $u'$  perturbation provided  $\bar{F} < \bar{F}_{CR}$ . Since the energy is a Lyapunov function for [7], trajectories off the  $\bar{u}$ -axis converge to either of the critical points  $(\bar{u}_*, \pm u'_{CR})$  with  $\bar{u}_*$  the marginally stable value; thus we can expect more turbulent behavior in the conceptual stochastic models with  $K \geq 2$  as the forcing  $\bar{F}$  increases in magnitude through negative values,  $\bar{F}$  with  $\bar{F} \leq \bar{F}_{CR} = -0.0067$ . A similar scenario occurs for the case with negative damping in [7] for  $\bar{F} \leq -0.0545$  with a single critical point along the  $\bar{u}$ -axis which is unstable to perturbations in  $u'$  with two critical points  $(\bar{u}_*, \pm u'_{CR})$ ,  $u'_{CR} \neq 0$ , which are also unstable because  $-\bar{d} - 3\bar{\alpha}\bar{u}_*^2 > 0$ ; in this case with all three equilibrium points unstable, trajectories off the  $\bar{u}$ -axis necessarily converge to periodic orbits encircling the critical points  $(\bar{u}_*, \pm u'_{CR})$  and frequently visit values of  $\bar{u}$  with instability in the  $u'$  dynamics. We also anticipate different behavior for  $\bar{F} > -0.0545$  since a stable critical point appears at  $\bar{u} = 0.8329$  for this and larger values of  $\bar{F}$ . See the tables in the supplementary material.


## Numerical Experiments for K=5 in the Conceptual Model

Here we use simple numerical experiments to demonstrate that the six dimensional conceptual model in [4] with  $K = 5$  has all the statistical features listed in (A)-(D) including intermittency of the small scales. The parameters,  $\bar{d}$ ,  $\bar{\alpha}$ , and  $\gamma = 1.5$  have already been discussed in [11]. The damping coefficients  $d_k$  are a mixture of uniform and scale selective damping with  $d_k = 1 + 0.02k^2$  for  $k = 1, 2, \dots, 5$  so that the smaller scales are damped more rapidly; the noise level set by  $\sigma_k$  for the  $k$ -th mode is determined by

$$\frac{\sigma_k^2}{d_k} = \frac{0.004}{(1+k)^{5/3}}, \quad k = 1, 2, \dots, 5 \quad [12]$$

so that a  $-5/3$  spectrum is calibrated to occur for these modes provided  $\bar{u} \equiv 0$  in the equations for  $u'_k$  [8]. This specifies all parameters in the conceptual model for turbulence used here. For all numerical simulations below and in the supplementary material, the Euler-Maruyama method is used with a time step  $\Delta t = 5 \times 10^{-3}$  and the system is integrated for a long time  $T = 2 \times 10^5$  with the first  $t = 2 \times 10^3$  time data ignored for post processing the climatological statistics. In


all simulations the initial value is  $\bar{u} = 1.5$  with  $u'_k = 0$  for  $k = 1, 2, \dots, 5$ .



**Fig. 1.** Negative large scale damping : time series (left column) and pdfs (right column) of the turbulent signal  $u$ ,  $\bar{u}$  and  $u'_k$ ,  $k = 1, 2, \dots, 5$  with  $\bar{F} = -0.055$ . Note the logarithmic scale of pdfs in the y-axis. Dashed lines are Gaussian distributions with the same mean and variance.

First we consider the case with large scale instability for  $\bar{u}$  with negative damping,  $\bar{d} = -0.1$  and  $\bar{\alpha} = 0.05$  with the forcing value  $\bar{F} = -0.055$  motivated by the phase portrait analysis above. Fig. 1 depicts the pdfs for the total turbulent field  $u$ , the large-scale mean  $\bar{u}$ , and the turbulent fluctuations  $u'_k$ ,  $k = 1, 2, \dots, 5$  as well as a sample of the time series of each variable in the conceptual model; the pdfs are plotted with a logarithmic vertical coordinate in order to highlight fat tails of intermittency while the Gaussian distribution with the same variance is the parabola in the figure. The pdf for the overall turbulent field  $u$  in [2] is nearly Gaussian while the pdfs for the mean  $\bar{u}$  and the largest scale fluctuating mode,  $u'_1$ , are both slightly sub-Gaussian. The variable  $u'_2$  has a Gaussian tail while the variables  $u'_3, u'_4, u'_5$  all have significant fat tails which are a hallmark of intermittency; the time series for  $u'_3, u'_4, u'_5$  in Fig. 1 clearly display highly intermittent behavior of extreme values with the amplitude of  $u'_5$  occasionally spiking to the typical amplitude of  $u'_1$  even through the climatological variance of  $u'_3$  is nearly eight times smaller than that for  $u'_1$  (see Table 2 of supplementary material). The climatological mean value for  $\bar{u}$  is  $-0.6733 = \langle \bar{u} \rangle$  and  $\langle \bar{u} \rangle$  is very close to the marginal stability value  $\bar{u}_* = -0.6667 = -\frac{d}{\gamma}$  motivated from [7] while the standard deviation of  $\bar{u}$  is 0.1993 indicating

that the instability mechanism elucidated in [5] is operating on all modes and creating intermittency. The total energy of the mean flow  $\bar{u}$  exceeds that of the fluctuations,  $u'_k$ . The variables  $u'_k, k = 1, 2, \dots, 5$  have essentially zero means with variances 0.0446, 0.0174, 0.0049, 0.0014, and 0.0005 respectively with the correlation time for  $\bar{u}$  approximately 34 while those for  $u'_k, k = 1, 2, \dots, 5$  are decreasing with  $k$  and approximately 29, 16, 6, 4, and 3 respectively. These are all the features of anisotropic turbulence required from (A)-(D) and demonstrated in the conceptual dynamical models; furthermore all of these conditions occur in a robust fashion for  $\bar{F}$  increasing in magnitude with  $\bar{F} \leq -0.055$  and  $0.055 \leq |\bar{F}| \leq 0.1$ . All of the detailed data discussed above can be found in Tables 1-3 of the supplementary material. There is an evident role for the unstable damping of the large scales,  $\bar{d} = -0.1$  to increase the variance of  $\bar{u}$  with its mean near the marginally critical value  $\bar{u}_*$  so that the instability mechanism from [5] operates vigorously in the model and creates more variance in  $u'_k, k = 1, 2, \dots, 5$ . Thus, we expect the system with stable damping and the same values of  $\bar{F}$  with  $\bar{F} = -0.055$  to have less variance.



**Fig. 2.** Positive large scale damping : time series (left column) and pdfs (right column) of the turbulent signal  $u, \bar{u}$  and  $u'_k, k = 1, 2, \dots, 5$  with  $\bar{F} = -0.080$ . Note the logarithmic scale of pdfs in the y-axis. Dashed lines are Gaussian distributions with the same mean and variance.

We consider the case with positive large scale damping,  $\bar{d} = 0.01$ , for  $\bar{F} = -0.080$  and in Fig. 2 we show the pdfs of all variables as well as a piece of the time series of the turbulent signal  $u, \bar{u}$ , and  $u'_k, k = 1, 2, \dots, 5$ . The intermittency of the small scale modes with less variance is evident

in Fig. 2. The mean flow variable,  $\bar{u}$ , has the largest total energy with climatological mean  $\langle \bar{u} \rangle = -0.6853$  which is very close to the marginal critical values  $\bar{u}_* = -0.6667$  so the intermittent instability mechanism in [5] is operating once again. Both the variances and correlation times behave in a similar fashion as for the negative large scale damping case discussed above and as required in (A)-(D) so the conceptual model with positive large scale damping also is a qualitative dynamical model for anisotropic turbulence with all the features in (A)-(D). Furthermore, all of these features persist for  $\bar{F}$  with  $-0.055 \leq \bar{F} \leq -0.1$ ; the pdfs are all Gaussian with no fat tails for  $\bar{F}$  with sufficiently small absolute value such as  $\bar{F} = -0.01$  as shown in the supplementary material. As expected from our discussion of the unstable case; for fixed forcing with  $\bar{F} \leq -0.055$ , there is between a factor of two and three less variance in all variables in the positive large scale damping case compared with the negative large scale damping case. Documentation for all of the above claims is found in extensive tables in the supplementary material. For both cases cross correlation among the variables  $\bar{u}, u'_k, k = 1, 2, \dots, 5$  are negligible in the climatological mean state with values roughly less than the 5% level.

In the above paragraphs, we emphasized models with  $K = 5$  to mimic the many degrees of freedom in real anisotropic turbulence and their interaction with the mean flow. From a mathematical viewpoint, it is interesting to address the following: what is the lowest dimensional conceptual model with intermittency and satisfying all the requirements in (A)-(D)? Versions of the conceptual model with  $K = 2$  already exhibit intermittency in  $u'_2$  as well as all the other features required in (A)-(D) for both positive and negative damping as shown in the supplementary material. However, the two mode models with  $K = 1$  always exhibit either sub-Gaussian or at most Gaussian behavior in  $u'_1$  without intermittency as the noise level is varied in all of our numerical experiments.

## Concluding Discussion

Conceptual dynamical models for anisotropic turbulence have been introduced here which, despite their simplicity, capture key features of vastly more complicated systems. The conceptual dynamical models introduced here in [4] involve a large scale mean flow  $\bar{u}$  and turbulent fluctuations,  $u'_k, 1 \leq k \leq K$ , on variety of spatial scales and involve energy conserving wave-mean flow interactions as well as suitable degenerate stochastic forcing of the fluctuations  $u'_k$ . The models have a transparent mechanism where the mean flow,  $\bar{u}$ , can destabilize the  $k$ -th mode whenever  $d_k + \gamma\bar{u} < 0$ ; a phase plane analysis yields parameters and robust regimes of sufficiently strong large scale external forcing,  $\bar{F}$ , where the models have a climatological mean state  $\langle u \rangle$  which is nearly neutrally stable in the sense that  $d_1 + \gamma\langle \bar{u} \rangle \cong 0$  so that fluctuations in the mean  $\bar{u}$  often introduce intermittent instability. Numerical experiments with a six-dimensional version of the model summarized here and in the supplementary material confirm that it captures key statistical features of vastly more complex anisotropic turbulent systems. These include chaotic statistical behavior of the mean flow,  $\bar{u}$ , with a sub-Gaussian pdf for its fluctuations while the turbulent fluctuations,  $u'_k, 1 \leq k \leq 5$ , have decreasing energy and correlation times as  $k$  increases with nearly Gaussian pdfs for the large scale fluctuations and fat-tailed non-Gaussian pdfs for the smaller scale fluctuations; this last feature allows for intermittency of the small scale fluctuations where turbulent modes with small variance can have relatively frequent large amplitude extreme events which directly impact the mean flow,  $\bar{u}$ . Remarkably, vastly more complex realistic turbulent systems often exhibit such marginal

critical behavior on average [4]. As mentioned above [1, 2], we can regard  $\bar{u}, u'_k$  for  $1 \leq k \leq K$  as defining turbulent fluctuations at a grid point in a vastly more complex spatially extended system. There are straightforward generalization of the conceptual model to allow for many large scale grid points,  $\bar{u}_j, j = 1, 2, \dots, J$  with associated turbulent fluctuations  $u'_{j,k}, 1 \leq k \leq K$  satisfying a coupled system of equations on the large scales,

$$\begin{aligned} \frac{d\bar{u}_j}{dt} &= L\bar{u}_j + \gamma \sum_{k=1}^K (u'_{j,k})^2 - \bar{d}\bar{u}_j - \bar{\alpha}\bar{u}^3 + \bar{F}_j \\ \frac{du'_{j,k}}{dt} &= -(d_k + \gamma\bar{u}_{j,k})u'_{j,k} + \sigma_k \dot{W}_{j,k} \end{aligned} \quad [13]$$

1. Hinze J (1959) *Turbulence : An introduction to its mechanisms and theory* (McGraw-Hill New York)
2. Townsewd AA (1976) *The Structure of Turbulent Shear Flow* (Cambridge Univ. Press, Cambridge MA)
3. Frisch U (1995) *Turbulence* (Cambridge Univ. Press, New York)
4. Vallis GK (2006) *Atmospheric and oceanic fluid dynamics: fundamentals and large-scale circulation* (Cambridge Univ. Press, Cambridge MA)
5. Salmon R (1998) *Lectures on Geophysical Fluid Dynamics* (Oxford Univ. Press, New York)
6. Majda AJ, Grooms I (2014) New perspectives on superparameterization for geophysical turbulence, *J. Comp. Phys.* ISSN 0021-9991 doi:10.1016/j.jcp.2013.09.014.
7. Holmes P, Lumley JL, Berkooz G (1996) *Turbulence, Coherent Structures, Dynamical Systems and Symmetry* (Cambridge Univ. Press, New York)
8. Majda AJ, Harlim J (2012) *Filtering Complex Turbulent Systems* (Cambridge Univ. Press, New York)
9. Mattingly JC, Stuart AM, Higham DJ (2002) Ergodicity for SDEs and approximations : locally Lipschitz vector fields and degenerate noise, *Stoch. Proc. Appl.*, 101(2):185–232
10. Majda AJ, Grote MJ (2009) Mathematical test models for superparameterization in anisotropic turbulence, *Proc. Natl. Acad. Sci. USA* 106(14):5470–5474
11. Majda AJ, Branicki M (2012) Lessons in uncertainty quantification for turbulent dynamical systems, *Disc. and Cont. Dyn. Sys.* 32(9):3133–3221

where  $L$  can be a linear or nonlinear operator coupling the  $\bar{u}_j$ . The conceptual models in [13] are nonlinear generalizations with transparent physical mechanisms of those introduced to study stochastic superparameterization in anisotropic turbulence [10, 6]. Besides their role as qualitative analogue models of vastly more complicated anisotropic turbulence, the conceptual dynamical models introduced here are potentially useful as a simplified test bed for algorithms and strategies for prediction, uncertainty quantification [11], and data assimilation [8] in vastly more complex anisotropic turbulent systems.

**ACKNOWLEDGMENTS.** The research of A.J.M. is partially supported by Office of Naval Research (ONR) Grants, ONR-Departmental Research Initiative N0014-10-1-0554, ONR N0014-11-1-0306, and ONR Multidisciplinary University Research Initiative 25-74200-F7112. Y.L. is supported as a postdoctoral research fellow on this last award.