

Tiling the plane with equilateral triangles

János Pach* Gábor Tardos†

Abstract

Let \mathcal{T} be a tiling of the plane with equilateral triangles no two of which share a side. We prove that if the side lengths of the triangles are bounded from below by a positive constant, then \mathcal{T} is periodic and it consists of translates of only at most three different triangles. As a corollary, we prove a theorem of Scherer and answer a question of Nandakumar. The same result has been obtained independently by Richter and Wirth.

1 Introduction

A collection of compact convex sets with nonempty interiors is said to *tile* a region R if no two of them share an interior point and their union is equal to R .

In 1934, during his first visit to Trinity College, Cambridge, P. Erdős [19] raised the following question: Is it possible to tile a unit square with finitely many smaller squares, no two of which are congruent? It has taken a few years before four promising students at Trinity College, R. L. Brooks, C. A. B. Smith, A. H. Stone, and W. T. Tutte [4], all of whom became prominent mathematicians, managed to answer Erdős's question in the affirmative. They discovered and explored an intimate relationship between tilings and flows in networks, and constructed several tilings with the required properties. In any such tiling \mathcal{T} of a unit square S , there is a unique *smallest* square. This square cannot share a side with the boundary of S , otherwise one of its neighbors would be congruent to it. Scaling up \mathcal{T} so that the side length of its smallest square becomes 1, and tiling this unit square with a congruent copy of \mathcal{T} , we obtain a tiling $\mathcal{T}' \supset \mathcal{T}$ of a larger square with pairwise noncongruent squares. Repeating this procedure infinitely many times, we obtain the following.

Corollary. *The plane can be tiled by infinitely many pairwise noncongruent squares whose side lengths are bounded from below by a positive constant.*

*EPFL, Lausanne and Rényi Institute, Budapest. Supported by Swiss National Science Foundation Grants 200020-162884 and 200021-165977. E-mail: pach@cims.nyu.edu.

†Rényi Institute, Budapest. Supported by the Cryptography “Lendület” project of the Hungarian Academy of Sciences and by the National Research, Development and Innovation Office, NKFIH, projects K-116769 and SNN-117879.

Tutte [20] extended the technique developed in [4] to tilings of an equilateral triangle with equilateral triangles. In particular, he proved that no such tiling exists with pairwise noncongruent triangles. However, this does not answer the corresponding question for tilings of the entire plane, which was raised by R. Nandakumar [13] (June 14, 2016).

Problem 1. (Nandakumar) *Is it possible to tile the plane with pairwise noncongruent equilateral triangles whose side lengths are bounded from below by a positive constant?*

In fact, we do not even know whether there exists a tiling of the plane with pairwise noncongruent equilateral triangles if we replace the condition that they cannot be arbitrarily small by the weaker assumption that the tiling is *locally finite*, that is, every bounded region intersects only finitely many triangles. Without assuming local finiteness, it is easy to find a tiling of the “punctured” plane and with a little more work of the whole plane; see [16].

Proposition 2. (Klaassen [9]) *There exists a tiling of the plane minus a single point with equilateral triangles of pairwise distinct sizes.*

Proof: Let α be the only real root of the polynomial $x^3 + x^2 - 1$. Let P and A be distinct points in the Euclidean plane. Let ϕ be the similarity transformation of the plane that has P as its fixed point, rotates the plane with angle $\pi/3$ and shrinks the distances with factor α . Let T_0 be the equilateral triangle with vertices A , $\phi(A)$ and a third vertex in the half plane bounded by the line $A\phi(A)$ and containing P .

The equilateral triangles $T_i = \phi^i(T_0)$, where i ranges over all integers, form a tiling of the plane minus the point P , and the sizes of these triangles are pairwise distinct. \square




Figure 1: A tiling with no two triangles sharing a side.

The aim of this note is to answer Nandakumar's above mentioned question (Problem 1).

Theorem 3. *Let \mathcal{T} be a tiling of the plane with equilateral triangles such that the side lengths of the triangles are bounded from below by a positive constant and no two triangles share a side. Then the triangles in \mathcal{T} have at most three different side lengths, a , b and c with $a = b + c$ (where c may be equal to b), and the tiling is periodic.*

Corollary 4. *There is no tiling of the plane with pairwise noncongruent equilateral triangles whose side lengths are bounded from below by a positive constant.*

The weaker statement that there is no tiling satisfying the conditions of Corollary 4, in which there is a *smallest* triangle, was proved by Scherer [18].

Nandakumar [12, 13], who teaches undergraduates at a local college in Cochin, India, raised several other interesting questions about tilings. They have triggered a lot of research in geometry and topology; see [1, 2, 5, 8, 10, 11, 14].

After completing our note, it has come to our attention that Theorem 3 has also been proved independently and probably slightly earlier by Richter and Wirth [17]. Their proof is somewhat different from ours, but in the last step we both use martingales. We are grateful to Richter and Wirth to their helpful remarks on our manuscript.

2 Proof of Theorem 3

Let us fix tiling \mathcal{T} satisfying the requirements of Theorem 3. We will prove the statement of the theorem through a sequence of lemmas about \mathcal{T} . First, we agree about the terminology.

We call an edge of a triangle $T \in \mathcal{T}$ *subdivided* if some interior point of this edge is a vertex of another triangle $T' \in \mathcal{T}$. Otherwise, it is called *uncut*. Note that a vertex in the tiling is the interior point of at most one edge and in that case it is the vertex of exactly three triangles.

We say that the edge AB of a triangle *continues at A* if A is the interior point of an edge e of another triangle of our tiling and e also contains some interior points of the edge AB . Clearly, at most one of the two edges of a triangle meeting at a vertex can continue there. By our assumptions, the same segment cannot be the edge of two triangles in the tiling. Therefore, every uncut edge must continue at least at one of its endpoints.

We will use the following simple observation several times.

Lemma 5. *If an edge AB is subdivided and does not continue at A , then there is a unique triangle ADE in the tiling such that D is an interior point of AB . The edge AD is uncut.* \square

We call a triangle of the tiling *small* if all three of its sides are uncut. We call it *large* if all three sides are subdivided. We call it *improper* if it has an uncut side and also a side that does not continue in either direction.




Figure 2: Illustration to Lemma 6.

Lemma 6. *For any improper triangle $T \in \mathcal{T}$, there are two other triangles, $U, V \in \mathcal{T}$ such that U is improper and the sum of the side lengths of U and V does not exceed the side length of T .*

Proof: let $T = ABC$ and assume that the edge BC is uncut and AB does not continue at either endpoint. As we noted, AB must be subdivided in this case and, by Lemma 5, we can find distinct triangles ADE and $D'BF$ such that D and D' are (not necessarily different) interior points of the segment AB . See Figure 1. We clearly have $AD + D'B \leq AB$ and we are done, unless neither of these triangles is improper. Assume that they are not improper. Their edges AD and $D'B$ are uncut, so all their edges have to continue in at least one direction. The edge AD continues at D , so DE cannot continue there, so it must continue at E . Therefore, the edge AE must continue at A . Similarly, FB must continue at B . As BC is uncut it must continue at one of its endpoints. As FB continues at B , the edge BC cannot continue there, so it must continue at C . The edge AC does not continue in either direction, since AE continues at A and BC continues at C .

As in the previous paragraph, we can find the triangles AGH and $G'CI$ in the tiling with G, G' interior points of the segment AC and we are done, unless neither of them is improper. But, as we have seen earlier, in this case IC would continue at C , contradicting the fact BC continues there. This contradiction completes the proof of the lemma. \square

Lemma 7. *There is no improper triangle in the tiling.*

Proof: Suppose T_0 is improper. By Lemma 6, for every $i \geq 1$ we can recursively find triangles T_i and $V_i \in \mathcal{T}$ such that T_i is improper and the sum of the side lengths of T_i and V_i does not

exceed the side length of T_{i-1} . Since the sum of the side lengths of all triangles V_i is at most the side length of T_0 , this sum must be convergent, contradicting our assumption that the sizes of all triangles in \mathcal{T} are separated from zero. The contradiction proves the lemma. \square

Lemma 8. *If the edge AB of a triangle ABC in \mathcal{T} continues at B , then BC is uncut.*

Proof: As AB continues at B , BC does not continue there, so if it is subdivided, then we have a triangle BDE such that D is an interior point of BC (Lemma 5). The edge BE does not continue at B (AB continues there), and DE does not continue at D (BD continues there). At most one of BE and DE can continue at E , so the triangle ADE must have at least one edge that does not continue in either direction. It also has an uncut edge, BD , so it is improper. The contradiction with Lemma 7 proves Lemma 8. \square

Lemma 9. *Every triangle in \mathcal{T} is either large or small. The sides of the large triangles do not continue in either direction, and each of them contains in its interior precisely one vertex.*

Proof: Assume first that the triangle ABC has an edge that continues in at least one direction, say AB continues at B . By Lemma 8, BC is uncut, so it must also continue in one direction. It cannot continue at B , hence it continues at C . By the same lemma, AC must be uncut and it must continue at A . Finally, using Lemma 8 again, we obtain that AB is uncut and, therefore, ABC is a small triangle.

Assume next that none of the edges of ABC continues in either direction. Then all edges must be subdivided and ABC is a large triangle. It remains to prove that all three edges contain exactly one subdividing vertex. Indeed, if AB contains more than one, then by an observation similar to Lemma 5, we have a triangle DEF where both D and E are interior vertices of the edge AB . In this case, DF and EF do not continue at D and E , respectively, (DE continues at both vertices), and at most one of DF and EF can continue at F . So, DEF has an edge that does not continue in either direction. It also contains an uncut edge DE , so it is improper. The contradiction with Lemma 7 completes the proof of Lemma 9. \square

According to Lemma 9, the combinatorial structure of the triangulation \mathcal{T} is completely determined. A small part of the tiling is depicted in Figure 2. Each vertex of the triangulation is the vertex of exactly one large triangle and exactly two small triangles, and it is contained in exactly one further triangle: it is an interior vertex of an edge of a large triangle.

The following observation is immediate, using the fact that no triangles in the tiling are improper.

Lemma 10. *Let $T = ABC$ be a large triangle in the tiling and T_1, T_2 and T_3 be the large triangles containing A, B and C , respectively, in the interior of one of their edges. Then the side length of T is the average of the side lengths of T_1, T_2 and T_3 . \square*

Proof of Theorem 3: Consider two large triangles T and T' . We claim that they must be congruent.

Assume for contradiction that the side length a of T is smaller than the side length a' of T' . Start a random walk at $T_0 = T$. After visiting the large triangle T_i , choose the next large




Figure 3: The combinatorial structure of \mathcal{T} .

triangle T_{i+1} uniformly at random from among the three triangles that contain a vertex of T_i in the interior of one of its edges. Stop the random walk at the first time when $T_i = T'$ holds.

According to a well known extension of Pólya's recurrence theorem [15], this random walk is recurrent; see [7], Theorem 8.2 on p. 138. Thus, choosing N large enough, we can make sure that after N steps it reaches T' with probability at least a/a' . We stop the process at step N , even if T' has not been reached.

The side lengths of T_i form a martingale by Lemma 10. So the expected side length at the end of the process is exactly the side length a of $T_0 = T$. At the end, this side length is a' with probability at least a/a' and it is always positive, so we have $a > (a/a')a'$. This contradiction proves that all large triangles must be of the same size.

In view of Lemma 8, every side of a large triangle is subdivided into two segments belonging to two small triangles. Since the structure of the tiling is uniquely determined, the small triangles must fall into at most two congruence classes, and the tiling is periodic (in fact, it is the union of three latticelike arrangements; see [6]). \square

References

- [1] I. Bárány, P. Blagojević, and A. Szűcs: Equipartitioning by a convex 3-fan, *Advances in Math.* **223** (2010), 579–593.

- [2] P. Blagojević and G. Ziegler: Convex equipartitions via equivariant obstruction theory, *Israel J. Math.* **200** (2014), no. 1, 49–77.
- [3] P. Brass, W. Moser, and J. Pach: *Research Problems in Discrete Geometry*, Springer, New York, 2005.
- [4] R. L. Brooks, C. A. B. Smith, A. H. Stone, and W. T. Tutte: The dissection of rectangles into squares, *Duke Math. J.* **7** (1940), 312–340.
- [5] D. Frettlöh: Noncongruent equidissections of the plane, [arXiv:1603.09132](https://arxiv.org/abs/1603.09132).
- [6] P. M. Gruber: *Convex and Discrete Geometry. Grundlehren der Mathematischen Wissenschaften* Vol. **336**, Springer, Berlin, 2007.
- [7] O. Kallenberg: Foundations of Modern Probability, in series: *Probability and its Applications*, Springer, New York, 1977.
- [8] R. Karasev, A. Hubard, and B. Aronov: Convex equipartitions: the spicy chicken theorem, *Geom. Dedicata* **170** (2014), 263–279.
- [9] B. Klaassen: Infinite perfekte Dreieckszerlegungen auch für gleichseitige Dreiecke. (German) [Infinite perfect dissection of triangles including equilateral triangles] *Elem. Math.* **50** (1995), no. 3, 116–121.
- [10] A. Kupavskii, J. Pach, and G. Tardos: Tilings with noncongruent triangles, preprint, [arXiv:1711.04504](https://arxiv.org/abs/1711.04504).
- [11] A. Kupavskii, J. Pach, and G. Tardos: Tilings of the plane with unit area triangles of bounded diameter, *Acta Math. Hungar.*, to appear.
- [12] R. Nandakumar, Fair partitions. Blog entry, <http://nandacumar.blogspot.de>, September 28, 2006.
- [13] R. Nandakumar: Filling the plane with non-congruent pieces, blog entries, December 2014, January 2015, June 2016, <http://nandacumar.blogspot.in>.
- [14] R. Nandakumar and N. Ramana Rao: Fair partitions of polygons: An elementary introduction, *Proc. Indian Academy of Sciences (Mathematical Sciences)* **122** (2012), 459–467.
- [15] G. Pólya: Über eine Aufgabe betreffend die Irrfahrt im Strassennetz, *Math. Ann.* **84** (1921), 149–160.
- [16] Ch. Richter: Tiling by incongruent equilateral triangles without requiring local finiteness, *Elem. Math.* **67** (2012), no. 4, 157–163.
- [17] Ch. Richter and M. Wirth: Tilings of convex sets by mutually incongruent equilateral triangles contain arbitrarily small tiles, manuscript arxiv.org/abs/1711.08903.

- [18] K. Scherer: The impossibility of a tessellation of the plane into equilateral triangles whose sidelengths are mutually different, one of them being minimal, *Elem. Math.* **38** (1983), no. 1, 1–4.
- [19] B. Schechter: *My Brain is Open*, Simon & Schuster, New York, 1998.
- [20] W. T. Tutte: The dissection of equilateral triangles into equilateral triangles, *Proc. Cambridge Philos. Soc.* **44** (1948), 463–482.